

PRODUCT MONOGRAPH

 **VICTRELIS TRIPLE™**

VICTRELIS™
boceprevir capsules, 200 mg
Hepatitis C Virus (HCV) Protease Inhibitor

PLUS

PEGETRON®
ribavirin capsules, 200 mg
Antiviral Agent

plus

peginterferon alfa-2b powder for solution in REDIPEN® single dose delivery system:
80 mcg/0.5 mL
100 mcg/0.5 mL
120 mcg/0.5 mL
150 mcg/0.5 mL, when reconstituted
Biological Response Modifier

NOTE: THESE PRODUCTS ARE INTENDED ONLY FOR USE AS DESCRIBED. THE INDIVIDUAL PRODUCTS CONTAINED IN VICTRELIS TRIPLE™ SHOULD NOT BE USED ALONE OR IN COMBINATION FOR OTHER PURPOSES. THE INFORMATION DESCRIBED IN THIS PRODUCT MONOGRAPH CONCERNS ONLY THE USE OF THESE PRODUCTS. FOR INFORMATION ON THE USE OF THE INDIVIDUAL COMPONENTS WHEN DISPENSED AS INDIVIDUAL MEDICATIONS OUTSIDE THIS COMBINED USE FOR THE TREATMENT OF CHRONIC HEPATITIS C (CHC), THE RESPECTIVE PRODUCT MONOGRAPHS FOR THESE PRODUCTS SHOULD BE CONSULTED.

Merck Canada Inc.

16750, route Transcanadienne
Kirkland, QC Canada H9H 4M7
<http://www.merck.ca>

Date of Preparation:
November 4, 2011

Submission Control No: 142814

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION..... 3
SUMMARY PRODUCT INFORMATION 3
DESCRIPTION..... 3
INDICATIONS AND CLINICAL USE..... 3
CONTRAINDICATIONS 4
WARNINGS AND PRECAUTIONS..... 6
ADVERSE REACTIONS..... 16
DRUG INTERACTIONS 26
DOSAGE AND ADMINISTRATION..... 36
OVERDOSAGE 43
ACTION AND CLINICAL PHARMACOLOGY 44
STORAGE AND STABILITY..... 50
SPECIAL HANDLING INSTRUCTIONS 51
DOSAGE FORMS, COMPOSITION AND PACKAGING 51

PART II : SCIENTIFIC INFORMATION 53
PHARMACEUTICAL INFORMATION..... 53
CLINICAL TRIALS..... 55
DETAILED PHARMACOLOGY 65
MICROBIOLOGY 76
TOXICOLOGY 79
REFERENCES 86

PART III: CONSUMER INFORMATION..... 88

VICTRELIS TRIPLE™

boceprevir, ribavirin plus peginterferon alfa-2b

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Non-medicinal Ingredients ^a
boceprevir: oral	capsules / 200 mg	lactose monohydrate
ribavirin: oral	capsules / 200 mg	lactose monohydrate
peginterferon alfa-2b: subcutaneous	powder for solution in REDIPEN [®] single dose delivery system / 80 mcg/0.5 mL 100 mcg/0.5 mL 120 mcg/0.5 mL 150 mcg/0.5 mL	none

a: For a complete listing see **DOSAGE FORMS, COMPOSITION AND PACKAGING** section.

DESCRIPTION

VICTRELIS TRIPLE™ is a combination of VICTRELIS™ (boceprevir (BOC)) and PEGETRON[®] (ribavirin (RBV) and peginterferon alfa-2b (PegIFN α 2b)), specifically BOC capsules, RBV capsules and PegIFN α 2b powder for solution.

VICTRELIS™

Boceprevir capsules: Boceprevir is an inhibitor of the hepatitis C virus (HCV) NS3/4A protease.

Boceprevir is manufactured as an approximately equal mixture of two diastereoisomers; the pharmacologically active SCH 534128 (S-isomer) and SCH 534129 (R-isomer).

PEGETRON[®]

Ribavirin capsules: Ribavirin is a synthetic nucleoside analog, which has shown *in vitro* activity against some, but not all, RNA and DNA viruses.

Peginterferon alfa-2b powder for solution: Peginterferon alfa-2b is a covalent conjugate of recombinant interferon alfa-2b with monomethoxy polyethylene glycol (PEG, with an average molecular weight of 12,000 daltons). Peginterferon alfa-2b is a biological response modifier.

INDICATIONS AND CLINICAL USE

VICTRELIS TRIPLE™ (boceprevir, ribavirin plus peginterferon alfa-2b) is indicated for: The treatment of Chronic Hepatitis C (CHC) genotype 1 infection in adult patients (18 years and older) with compensated liver disease, including cirrhosis, who are previously untreated or who have failed previous therapy.

This indication is based on evidence from two Phase 3 clinical studies conducted with VICTRELIS™ in combination with PegIFNα2b/RBV, SPRINT-2 in previously untreated patients and RESPOND-2 in previous treatment failures. Previous null responders (defined by a < 2.0 log₁₀ decline in HCV-RNA at Treatment Week (TW) 12) were excluded from the RESPOND-2 study. However, the results of the Phase 3 studies included a post-hoc analysis of a population similar to Null Responders (defined as poorly interferon responsive at TW 4 [$< 1.0 \log_{10}$ decline in HCV-RNA]) (see **CLINICAL TRIALS**).

Treatment with VICTRELIS TRIPLE™ should be initiated and monitored by a physician experienced in the management of CHC. Before initiating therapy, the following points should be considered:

- VICTRELIS TRIPLE™ efficacy has not been studied in patients who have failed previous therapy regimens which included VICTRELIS™ or other HCV NS3/4A Protease Inhibitors (PIs).
- Poorly interferon responsive patients who were treated with VICTRELIS TRIPLE™ achieved lower Sustained Virologic Response (SVR) and higher rate of resistance associated substitutions upon treatment failure compared to patients with a greater response to PEGETRON® (PegIFNα2b plus RBV) (see **CLINICAL TRIALS** and **MICROBIOLOGY, Resistance**).

Geriatrics (> 65 years of age)

Clinical studies of VICTRELIS TRIPLE™ included only a limited number of patients aged 65 and over. Consideration should be given to the decreased hepatic, renal or cardiac function and of concomitant disease or other drug therapy in elderly patients prior to prescribing VICTRELIS TRIPLE™ (see **WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION** and **ACTION AND CLINICAL PHARMACOLOGY**).

Pediatrics (< 18 years of age)

No data is available (see **WARNINGS AND PRECAUTIONS** and **ACTION AND CLINICAL PHARMACOLOGY**).

CONTRAINDICATIONS

VICTRELIS TRIPLE™ is contraindicated in:

Patients with known hypersensitivity to BOC, RBV and/or any interferons or any component of the capsules and/or injection. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section.

Patients with autoimmune hepatitis or a history of autoimmune disease.

Patients with hepatic decompensation (Child-Pugh score > 6 [class B and C]).

Pregnant women or by men whose female partners are pregnant. Extreme care must be taken to avoid pregnancy in female patients and in female partners of male patients taking VICTRELIS TRIPLE™ therapy. VICTRELIS TRIPLE™ therapy must not be initiated until a report of a negative pregnancy test has been obtained immediately prior to initiation of therapy. Women of childbearing potential and their male partners must not receive VICTRELIS TRIPLE™ therapy unless they are using effective contraception (two reliable forms, one for each partner) during treatment with VICTRELIS TRIPLE™ and for the 6-month post-therapy period (i.e., 15 half-lives for RBV clearance from the body). Significant teratogenic and/or embryocidal effects have been demonstrated for RBV in all animal species in which adequate studies have been conducted. These effects occurred at doses as low as one twentieth of the recommended human dose of RBV (see **WARNINGS AND PRECAUTIONS**).

Patients who have pre-existing severe psychiatric condition or a history of severe psychiatric disorder.

Patients who have pre-existing thyroid abnormalities for which thyroid function cannot be maintained in the normal range by medication.

Patients with severe renal dysfunction (creatinine clearance < 50 mL/min).

Patients who have epilepsy.

Co-administration of BOC with medicines that are highly dependent on CYP3A4/5 for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events (narrow therapeutic index). These drugs are listed in Table 1 (see **DRUG INTERACTIONS, Drug-Drug Interactions**).

Co-administration with medicines that are potent CYP3A4/5 inducers, where significantly reduced BOC plasma concentrations may be associated with reduced efficacy. These drugs are listed in Table 1 (see **DRUG INTERACTIONS, Drug-Drug Interactions**).

Table 1: Drugs that are contraindicated with boceprevir

Drug Class/Drug Name	Clinical Comment
Alpha 1-Adrenoreceptor Antagonists Alfuzosin	Increased alfuzosin concentrations can result in hypotension.
Antiarrhythmics amiodarone, propafenone, quinidine	Potential to produce serious and/or life-threatening Adverse Events (AEs).
Anticonvulsants carbamazepine, phenobarbital, phenytoin	May lead to loss of virologic response to VICTRELIS™.
Antimycobacterials Rifampin	May lead to loss of virologic response to VICTRELIS™.
Ergot Derivatives dihydroergotamine, ergonovine, ergotamine, methylergonovine	Potential for acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues.
Gastrointestinal Motility Agents cisapride ^a	Potential for cardiac arrhythmias.

Drug Class/Drug Name	Clinical Comment
Herbal Products St. John's Wort (<i>Hypericum perforatum</i>)	May lead to loss of virologic response to VICTRELIS™.
HMG-CoA Reductase Inhibitors lovastatin, simvastatin	Potential for myopathy, including rhabdomyolysis.
Neuroleptics Pimozide	Potential for cardiac arrhythmias.
Oral Contraceptives Drospirenone	Potential for hyperkalemia.
PDE-5 Enzyme Inhibitors sildenafil or tadalafil when used for the treatment of Pulmonary Arterial Hypertension (PAH)	Potential for PDE5 inhibitor-associated AEs, including visual abnormalities, hypotension, prolonged erection, and syncope.
Second Generation Antihistamines astemizole ^a , terfenadine ^a	Potential for cardiac arrhythmias.
Sedatives/Hypnotics midazolam (orally administered), triazolam (orally administered)	Potential for prolonged or increased sedation or respiratory depression.

a: Please note that cisapride, astemizole and terfenadine are no longer available on the Canadian market.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Alpha interferons, including PegIFN α 2b, cause or aggravate fatal or life-threatening neuropsychiatric, autoimmune, ischemic and infectious disorders. Patients should be monitored closely with periodic clinical and laboratory evaluations. Patients with persistently severe or worsening signs or symptoms of these conditions should be withdrawn from therapy. In many cases, but not all cases, these disorders resolve after stopping interferon therapy.

Life-threatening or fatal psychiatric effects, particularly severe depression, suicidal behavior (suicidal ideation, attempted suicide and suicide), psychosis including hallucinations, and aggressive behavior, sometimes directed towards others, have occurred in patients with and without previous psychiatric disorders during RBV plus PegIFN α 2b or RBV plus interferon alfa-2b treatment and follow-up (see **ADVERSE REACTIONS**). Other Central Nervous System (CNS) effects including confusion and alterations of mental status have been observed with alpha interferons, including PegIFN α 2b.

VICTRELIS TRIPLE™ therapy should be used with extreme caution in patients with a history of pre-existing psychiatric disorders who report a history of severe depression. If patients develop psychiatric or CNS problems, including clinical depression, it is recommended that the patients be carefully monitored during treatment and in the 6 month follow-up period, due to the potential seriousness of these undesirable effects. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician. If symptoms persist or worsen, discontinue VICTRELIS TRIPLE™ therapy.

General

VICTRELIS TRIPLE™ therapy should not be used in patients with severe, debilitating medical conditions.

Based on results of clinical trials, the use of RBV as monotherapy is not effective and RBV must not be used alone. The safety and efficacy of RBV has only been established in combination with PegIFN α 2b and in combination with PegIFN α 2b and VICTRELIS™.

VICTRELIS TRIPLE™ may cause moderate to severe adverse experiences requiring dose reduction or temporary interruption of RBV and/or PegIFN α 2b. Generally, medical management of these adverse experiences is accomplished with these modifications. Occasionally, discontinuation from further therapy is required. More stringent dose-modification guidelines are recommended for cardiac patients (see **DOSAGE AND ADMINISTRATION**).

Effects on Ability to Drive and Use Machines

No studies on the effects of VICTRELIS TRIPLE™ on the ability to drive and use machines have been performed. However, certain side effects that have been reported may affect some patients' ability to drive and use machines. Patients should be informed that fatigue, somnolence, confusion, dizziness, syncope, blood pressure fluctuations and blurred vision have been reported (see **ADVERSE REACTIONS, Clinical Trial Adverse Reactions**).

Carcinogenesis and Mutagenesis

Conventional carcinogenicity studies in rodents with low exposures compared to human exposure under therapeutic conditions (factor 0.1 in rats and 1 in mice) did not reveal tumorigenicity of RBV. In addition, in a 26 week carcinogenicity study using the heterozygous p53 (+/-) mouse model, RBV did not produce tumors at the maximally tolerated dose of 300 mg/kg (plasma exposure factor approximately 2.5 compared to human exposure). Ribavirin was non-carcinogenic when administered for 2 years to rats at doses up to 40 mg/kg (estimated human equivalent of 5.71 mg/kg based on body surface area adjustment for a 60 kg adult). While these studies do not suggest a carcinogenic potential of RBV in humans, this dose was less than the maximum tolerated dose, and therefore the 2-year study was not adequate to fully characterize the carcinogenic potential of RBV. Ribavirin should be considered a potential carcinogen.

Ribavirin is genotoxic and mutagenic in some *in vivo* and *in vitro* genotoxicity assays. Studies indicated that RBV was not oncogenic in mice at oral gavage doses up to 75 mg/kg/day, or in rats at oral gavage doses up to 40 mg/kg/day (see **TOXICOLOGY**).

See also **TOXICOLOGY** section for animal data.

Cardiovascular

Chest pain, hypertension, cardiac arrhythmia, cardiac ischemia, and myocardial infarction have been reported in patients with and without a history of cardiac disorder or abnormality in association with the use of alpha interferon therapies including PegIFN α 2b (see **ADVERSE REACTIONS**). VICTRELIS TRIPLE™ should not be administered to patients with a history of severe pre-existing cardiac disease including unstable or uncontrolled cardiac disease in the previous 6 months. As with other alpha interferons, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders who received PegIFN α 2b require close monitoring. It is recommended that patients who have pre-existing

cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of VICTRELIS TRIPLE™ therapy. Cardiomyopathy, that may be reversible upon discontinuation of alpha interferon, has been reported rarely in patients without prior evidence of cardiac disease.

A case of deep vein thrombosis has been reported from a clinical trial of retreatment of patients who failed previous treatment with interferon and RBV combination therapy.

Hydration

Adequate hydration must be maintained in patients undergoing therapy with VICTRELIS TRIPLE™, since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

Cerebrovascular Disorders

Ischemic and hemorrhagic cerebrovascular events have been observed in patients treated with interferon alpha-based therapies, including PegIFN α 2b. Events occurred in patients with few or no reported risk factors for stroke, including patients less than 45 years of age. Because these are spontaneous reports, estimates of frequency cannot be made and a causal relationship between interferon alpha-based therapies and these events is difficult to establish.

Dental and Periodontal Disorders

Dental and periodontal disorders have been reported in patients who received RBV and peginterferon combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of RBV and PegIFN α 2b. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

Ear/Nose/Throat

Hearing disorder and hearing loss have been reported with the use of alpha interferons, including PegIFN α 2b.

Endocrine and Metabolism

Diabetes Mellitus and Hyperglycemia

As with other alpha interferons, diabetes mellitus and hyperglycemia have been observed in patients treated with PegIFN α 2b. Symptomatic patients should have their blood glucose measured and followed up accordingly. Patients with diabetes mellitus may require adjustment of their anti-diabetic regimen.

Hypertriglyceridemia

Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed with PegIFN α 2b. Monitoring of lipid levels is, therefore, recommended.

Thyroid Changes

Infrequently, patients treated for chronic hepatitis C with alpha interferons, including PegIFN α 2b, have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. After discontinuation of therapy, thyroid dysfunction may or may not be reversed. Determine thyroid-stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, VICTRELIS TRIPLE™ treatment may be initiated or continued only if TSH levels can be maintained in the normal range by medication.

Gastrointestinal

Colitis

As seen with other alpha interferons, ulcerative and ischemic colitis, sometimes serious, have been observed within 12 weeks of starting PegIFN α 2b. VICTRELIS TRIPLE™ should be discontinued immediately if symptoms of colitis develop (typical manifestations include abdominal pain, bloody diarrhea and fever). The colitis usually resolves within 1 to 3 weeks of discontinuation of alpha interferon.

Hematologic

Anemia

Hemolytic anemia (hemoglobin levels to < 100 g/L) was observed in up to 28% of patients treated with RBV in combination with PegIFN α 2b in clinical trials. Anemia occurred within 1–4 weeks of initiation of RBV.

Although RBV has no direct cardiovascular effects, anemia associated with RBV may result in deterioration of cardiac function and/or exacerbation of the symptoms of coronary disease. Patients should be assessed before initiation of therapy and should be appropriately monitored during therapy. If there is any deterioration of cardiovascular status, therapy should be suspended or discontinued (see **DOSAGE AND ADMINISTRATION**). Because cardiac disease may be worsened by drug-induced anemia, patients with a history of significant or unstable cardiac disease should not be treated with VICTRELIS TRIPLE™. VICTRELIS TRIPLE™ therapy should not be used in patients with hemoglobinopathies (e.g., thalassemia, sickle-cell anemia).

The addition of VICTRELIS™ to PEGETRON® is associated with an additional decrease in serum hemoglobin concentrations (approximately 10 g/L) (see **ADVERSE REACTIONS, Clinical Trial Adverse Reactions, Anemia** and **ADVERSE REACTIONS, Abnormal Hematologic and Clinical Chemistry Findings**).

Neutropenia

Neutropenia is a known interferon-related AE. In Phase 2 and 3 clinical trials, the proportion of subjects with neutrophil counts of less than $0.5 \times 10^9/L$ was higher in the VICTRELIS TRIPLE™ arms (7%) compared to patients who received only PEGETRON® (4%) (see **ADVERSE REACTIONS, Abnormal Hematologic and Clinical Chemistry Findings, Neutrophils and Platelets**). In the key clinical trials, three patients experienced severe (2) or life-threatening (1)

infections within 14 days of Grade 3 or 4 neutropenia and two patients experienced life-threatening neutropenia while receiving VICTRELIS TRIPLE™.

Hepatic/Biliary/Pancreas

Hepatic Function

Any patient developing liver function abnormalities or hepatopathy during treatment should be monitored closely. As with treatment with any interferon, discontinue treatment with VICTRELIS TRIPLE™ in patients who develop prolongation of coagulation markers or other markers of hepatic function, which might indicate liver decompensation (see **DOSAGE AND ADMINISTRATION**). The safety and efficacy of PegIFN α 2b have not been evaluated in patients with severe hepatic dysfunction. Therefore, VICTRELIS TRIPLE™ must not be used for these patients. VICTRELIS TRIPLE™ therapy should be discontinued for any patient developing signs and symptoms of liver failure. Patients should be tested for the presence of HCV antibodies. Other causes of chronic hepatitis including autoimmune hepatitis should be excluded.

Pancreatitis

Pancreatitis, sometimes life-threatening, has occurred in patients treated with alpha interferons, including PegIFN α 2b. VICTRELIS TRIPLE™ therapy should be suspended if symptoms or signs of pancreatitis are observed. VICTRELIS TRIPLE™ should be discontinued in patients diagnosed with pancreatitis.

Immune

Acute Hypersensitivity

Acute hypersensitivity reactions, (e.g., angioedema, bronchoconstriction and anaphylaxis), have been observed rarely during alpha interferon therapy. If such a reaction develops during treatment with VICTRELIS TRIPLE™, discontinue treatment and institute appropriate medical therapy immediately. As with other alpha interferons, urticaria has been observed rarely during PegIFN α 2b therapy. Transient rashes do not necessitate interruption of treatment.

Autoimmune Disease

As with other alpha interferons, the development of autoantibodies has been reported during treatment with PegIFN α 2b. Clinical manifestations of autoimmune disease during interferon therapy may occur more frequently in patients predisposed to the development of autoimmune disorders.

Bone Marrow Toxicity

Alpha interferons, including PegIFN α 2b, may suppress bone marrow function which may result in severe cytopenias. As with other alpha interferons, PegIFN α 2b may be very rarely associated with aplastic anemia. PegIFN α 2b dosing should be reduced or discontinued in patients developing decreases in neutrophil or platelet counts (see **DOSAGE AND ADMINISTRATION, Dose modification**).

Fever

While fever may be associated with the “flu-like” syndrome reported commonly during interferon

therapy, other causes of persistent fever must be ruled out.

Immunological Effects

A number of immune-mediated dermatological reactions associated with the use of alpha interferons have been reported ranging from erythema multiforme to more severe but very rare occurrences of Stevens Johnson Syndrome and toxic epidermal necrolysis.

Neurologic

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses. While these effects are generally reversible upon discontinuation of therapy, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of PegIFN α 2b.

Ophthalmologic

Ocular Changes

As with other alpha interferons, ophthalmologic disorders, including retinopathies (including macular edema), retinal hemorrhages, retinal artery or vein obstruction, serous retinal detachment, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilledema have been reported in rare instances after treatment with PegIFN α 2b (see **ADVERSE REACTIONS**). These events have been reported after treatment for several months, but also have been reported after shorter treatment periods. All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Because these ocular events may occur in conjunction with other disease states, periodic ocular examinations during VICTRELIS TRIPLE™ therapy are recommended in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of VICTRELIS TRIPLE™ therapy should be considered in patients who develop new or worsening ophthalmological disorders.

Renal

Renal Function

It is recommended that renal function be evaluated in all patients prior to initiation of VICTRELIS TRIPLE™ therapy and that patients be monitored closely during treatment (see **DOSAGE AND ADMINISTRATION**). Increases in serum creatinine levels have been observed in patients with renal insufficiency treated with interferons, including PegIFN α 2b. Patients with impairment of renal function should be closely monitored for signs and symptoms of toxicity, including increases in serum creatinine, and, should have their weekly dose of PegIFN α 2b reduced if medically appropriate (see **DOSAGE AND ADMINISTRATION**). Patients with impaired renal function and/or those over the age of 50 should be more carefully monitored with respect to the development of anemia. Patients with severe renal dysfunction (creatinine clearance < 50 mL/min) must not be treated with VICTRELIS TRIPLE™, as the clearance of PegIFN α 2b is reduced in patients with significant renal impairment (see **ACTION AND CLINICAL PHARMACOLOGY** and **CONTRAINDICATIONS**). If serum creatinine rises to > 2 mg/dL, VICTRELIS TRIPLE™ must be discontinued.

Respiratory

Pulmonary Changes

As with other alpha interferons, pulmonary infiltrates, pneumonitis, pulmonary hypertension, and pneumonia, occasionally resulting in fatality, have been observed rarely in PegIFN α 2b treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely. If appropriate, discontinue VICTRELIS TRIPLE™ therapy. Prompt discontinuation of therapy and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events (AEs). These symptoms have been reported more frequently when *Shosaikoto* (also known as *Xiao-Chai-Hu-Tang*), a Chinese herbal medication, has been administered concomitantly with alpha interferons. VICTRELIS TRIPLE™ should not be administered to patients with chronic obstructive pulmonary disease (COPD).

Sensitivity/Resistance

Hepatitis C Virus Protease Monotherapy

Based on results of clinical studies, BOC must not be used alone due to the high probability of increased resistance without combination anti-HCV therapies (see **MICROBIOLOGY, Resistance**).

It is unknown what effect therapy with VICTRELIS TRIPLE™ will have on the activity of subsequently administered HCV NS3/4A PIs, including re-treatment with VICTRELIS TRIPLE™.

Sexual Function/Reproduction

Effects on Fertility

No human data on the effect of VICTRELIS TRIPLE™ on fertility are available. Available pharmacodynamic/toxicological data in rats have shown effects of BOC /metabolites on fertility, which have been reversible (female rats) and partially reversible (male rats) (see **TOXICOLOGY**).

No reproductive toxicology studies have been performed using PegIFN α 2b in combination with RBV. However, evidence provided below for interferon and RBV when administered alone indicates that both agents have adverse effects on reproduction. It should be assumed that the effects produced by either agent alone will also be caused by the combination of the two agents.

Interferons, including PegIFN α 2b, may impair fertility. In studies of interferon administration in non-human primates, menstrual cycle abnormalities have been observed. Decreases in serum estradiol and progesterone concentrations have been reported in women treated with human leukocyte interferon. The effects of interferon on male fertility have not been studied. Therefore, a possible effect on male fertility should be considered. The genotoxicity of PegIFN α 2b was evaluated in bacterial (Ames) and mammalian cell clastogenicity (HPBL) assays, and was negative in both assays.

VICTRELIS TRIPLE™ therapy should be used with caution in fertile men. In studies in mice to evaluate the time course and reversibility of RBV-induced testicular degeneration at doses of 15 to 150 mg/kg/day (estimated human equivalent of 1.25–12.5 mg/kg/day, based on body surface area adjustment for a 60 kg adult; 0.2–0.8 times the maximum human 24-hour dose of RBV) administered for 3 or 6 months, abnormalities in sperm occurred. Upon cessation of treatment, essentially total recovery from RBV-induced testicular toxicity was apparent within one or two spermatogenesis cycles.

Skin

Psoriatic Disease and Sarcoidosis

Use of alpha interferons, including PegIFN α 2b, has been associated with exacerbating pre-existing psoriatic disease and sarcoidosis. Use of VICTRELIS TRIPLE™ in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

Special Populations

Pregnant Women

No effects on fetal development have been observed in rats and rabbits at BOC AUC exposures approximately 11.8- and 2.0-fold higher, respectively, than those in humans at the recommended dose of 800 mg Three Times Daily (TID). BOC has been shown in animals to distribute across the placenta to fetal blood and tissues (see **TOXICOLOGY**).

Significant teratogenic and/or embryocidal potential have been demonstrated for RBV in all animal species in which adequate studies have been conducted, occurring at doses as low as one twentieth of the recommended human dose. Malformations of the skull, palate, eye, jaw, limbs, skeleton and gastrointestinal tract were noted. The incidence and severity of teratogenic effects increased with escalation of the RBV dose. Survival of fetuses and offspring was reduced. In conventional embryotoxicity/teratogenicity studies in rats and rabbits, observed no effect dose levels were well below those for proposed clinical use (0.3 mg/kg/day for both the rat and rabbit; approximately 0.06 times the recommended human 24-hour dose of RBV). No maternal toxicity or effects on offspring were observed in a peri/postnatal toxicity study in rats dosed orally at up to 1 mg/kg/day (estimated human equivalent dose of 0.17 mg/kg based on body surface area adjustment for a 60 kg adult; approximately 0.01 times the maximum recommended human 24-hour dose of RBV).

Interferon alfa-2b has been shown to have abortifacient effects in *Macaca mulatta* (rhesus monkeys) at 15 and 30 million IU/kg (estimated human equivalent of 5 and 10 million IU/kg, based on body surface area adjustment for a 60 kg adult). This same effect is expected with PegIFN α 2b. High doses of other forms of interferon alpha and beta are known to produce dose-related anovulatory and abortifacient effects in rhesus monkeys.

Because of these risks, VICTRELIS TRIPLE™ is contraindicated in pregnant women and men whose female partners are pregnant. Extreme care must be taken to avoid pregnancy in female patients and in female partners of male patients while taking this combination - both during treatment and for 6 months after the completion of all treatment. VICTRELIS TRIPLE™

therapy should not be started unless a female patient has a negative pregnancy test immediately prior to initiation of treatment and routine monthly pregnancy tests during treatment and for 6 months after all treatment has ended.

Systemic hormonal contraceptives may not be as effective in women while taking VICTRELIS TRIPLE™ (see **DRUG INTERACTIONS, Drug-Drug Interactions**). Therefore, two alternative effective methods of contraception, including intrauterine devices and barrier methods, should be used in women during treatment with VICTRELIS TRIPLE™.

There are no studies with VICTRELIS TRIPLE™ in pregnant women.

Treatment and Post-treatment: Potential Risk to the Fetus

Because of the potential human teratogenic effects of RBV, male patients must be advised to take every precaution to avoid risk of pregnancy for their female partners during treatment with VICTRELIS TRIPLE™ and for six months after treatment has been concluded (i.e., 15 half-lives for RBV clearance from the body). Ribavirin accumulates intracellularly and is cleared from the body very slowly. In animal studies, RBV produced changes in sperm at doses below the clinical dose. It is unknown whether the RBV that is contained in sperm will exert its known teratogenic effects upon fertilization of the ova. In a study in rats, it was concluded that dominant lethality was not induced by RBV at doses up to 200 mg/kg for 5 days (estimated human equivalent doses of 7.14 - 28.6 mg/kg, based on body surface area adjustment for a 60 kg adult; up to 1.7 times the maximum recommended human dose of RBV).

Women of childbearing potential and their male partners must not receive VICTRELIS TRIPLE™ therapy unless the patient and his/her partner are using effective contraceptive (two reliable forms, one for each partner) during the therapy period. In addition, effective contraception should be utilized for six months post-therapy based on a multiple dose half-life ($t_{1/2}$) of RBV of 12 days.

Male patients and their female partners must practice effective contraception (two reliable forms, one for each partner) during treatment with VICTRELIS TRIPLE™ and for the 6-month post-therapy period (e.g., 15 half-lives for RBV clearance from the body).

If pregnancy occurs in a patient or partner of a patient during treatment or during the six months after treatment cessation, the patient must be advised of the significant teratogenic risk of RBV Capsules to the fetus and physicians should report such cases by calling Merck Canada Inc., Medical Information Department at 1-800-567-2594.

Nursing Women

It is not known whether RBV and/or PegIFN α 2b are excreted in human milk.

Available pharmacodynamic/toxicological data in animals have shown excretion of BOC/metabolites in milk (see **TOXICOLOGY**). A risk to the newborns/infants cannot be excluded.

Because of the potential for serious adverse reactions from VICTRELIS TRIPLE™ in nursing infants, nursing must be discontinued prior to the start of VICTRELIS TRIPLE™ therapy.

Pediatrics (< 18 years of age)

The safety, efficacy and Pharmacokinetic (PK) profile of VICTRELIS TRIPLE™ in pediatric patients below 18 years of age have not yet been established (see **INDICATIONS AND CLINICAL USE**). VICTRELIS TRIPLE™ is not recommended for use in children and adolescents under the age of 18 years.

Geriatrics (> 65 years of age)

Clinical studies of BOC had limited data on patients aged 65 and over to determine whether they respond differently from younger patients. In general, consideration should be given to the potentially decreased hepatic, renal or cardiac function and concomitant disease or other drug therapy in elderly and younger patients (see **DETAILED PHARMACOLOGY**).

Human Immunodeficiency Virus (HIV) Co-infection

The safety and efficacy of VICTRELIS TRIPLE™ for the treatment of CHC genotype 1 infection have not been established in patients co-infected with HIV and HCV.

Hepatitis B Virus (HBV) Co-infection

The safety and efficacy of VICTRELIS TRIPLE™ for the treatment of CHC genotype 1 infection in patients co-infected with HBV and HCV have not been studied.

Organ Transplant Recipients

The safety and efficacy of VICTRELIS TRIPLE™ for the treatment of CHC genotype 1 infection in liver or other organ transplant recipients have not been studied.

Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported but a causal association with interferon alpha therapy has not been established.

Monitoring and Laboratory Tests

Standard hematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of VICTRELIS TRIPLE™ therapy are:

- Hemoglobin: ≥ 120 g/L (females), ≥ 130 g/L (males)
- Platelets: $\geq 100 \times 10^9$ /L
- Neutrophil Count: $\geq 1.5 \times 10^9$ /L
- TSH Levels: must be within normal limits

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

Uric acid may increase due to hemolysis with RBV Capsules use; therefore, the potential for development of gout must be carefully monitored in predisposed patients.

Complete blood counts (with white blood cell differential counts) should be obtained before treatment, at TWs 2, 4, 8 and 12 and should be closely monitored at other time points as considered clinically appropriate. If serum hemoglobin is < 100 g/L, a decrease in dose or interruption of RBV may be warranted (see **ADVERSE REACTIONS, Clinical Trial Adverse Drug Reactions, Anemia**). Decreases in the neutrophil counts may require dose reduction or discontinuation of PegIFN α /RBV (see **ADVERSE REACTIONS, Abnormal Hematologic and Clinical Chemistry Findings**).

HCV-RNA levels should be monitored at TWs 8, 12, and 24, at the End of Treatment (EOT), during treatment follow-up, and for other time points as clinically indicated. In previously untreated subjects without cirrhosis, monitoring of HCV-RNA levels at TW 4 is recommended to determine interferon responsiveness. Use of a sensitive real-time reverse-transcription polymerase chain reaction (RT-PCR) assay for monitoring HCV-RNA levels during treatment is recommended. The assay should have a lower limit of HCV-RNA quantification of equal to or less than 25 IU/mL, and a limit of HCV-RNA detection of approximately 10–15 IU/mL. For the purposes of assessing Response-Guided Therapy (RGT) milestones, a confirmed “detectable but below limit of quantification” HCV-RNA result should not be considered equivalent to an “undetectable” HCV-RNA result.

ADVERSE REACTIONS

Overview

Potentially serious adverse reactions have been reported with VICTRELIS TRIPLE™ in controlled clinical trials (see **WARNINGS AND PRECAUTIONS** and **ADVERSE REACTIONS, General Safety Information**).

The most commonly reported adverse reactions ($\geq 25\%$) with PEGETRON® alone were fatigue, fever, headache, rigors, myalgia, insomnia, injection site inflammation, injection site reaction, “flu-like” symptoms, weight decreased, anorexia, nausea, arthralgia, myalgia, depression, irritability, dyspnea, alopecia, pruritus, anemia, and neutropenia. The most frequently reported adverse reactions were mostly mild to moderate in severity and were manageable without the need for modification of doses or discontinuation of therapy.

The most commonly reported adverse reactions in clinical trials with VICTRELIS TRIPLE™ were similar across all study arms. The most commonly reported adverse reactions (incidence > 35%) considered by investigators to be causally related to VICTRELIS TRIPLE™ in adult patients in clinical trials were fatigue, anemia, nausea, headache, and dysgeusia.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be

compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

VICTRELIS™

The safety of VICTRELIS TRIPLE™ (combination of VICTRELIS™ 800 mg TID with PEGETRON®) was assessed in 2,095 patients with CHC in one Phase 2, open-label trial and two Phase 3, randomized, double-blinded, placebo-controlled clinical trials. The Phase 2 study SPRINT-1 (P03523), evaluated VICTRELIS TRIPLE™ with or without a four-week lead-in period with PEGETRON® compared to PEGETRON® alone in patients who were previously untreated. The Phase 3 studies SPRINT-2 (P05216 - patients who were previously untreated) and RESPOND-2 (P05101 - patients who had failed previous therapy), evaluated the use of VICTRELIS TRIPLE™ with a four-week lead-in period with PEGETRON® alone compared to PEGETRON® alone (see **CLINICAL TRIALS**). The population studied had a mean age of 49 years (2% of patients were > 65 years of age), 61% were male, 82% were White and 15% Black, and 7% of the population had cirrhosis (based on liver histology). In the pooled studies, the median exposure was 201 days in patients who received VICTRELIS TRIPLE™ and 198 days in patients who received PEGETRON® alone.

During the four-week lead-in period with PEGETRON® alone, 2% (28/1,263) patients in the VICTRELIS™-containing arms experienced adverse reactions leading to discontinuation of treatment. During the entire course of treatment, the proportion of patients who discontinued treatment due to adverse reactions was 13% for patients who received VICTRELIS TRIPLE™ and 12% for patients who received PEGETRON® alone. Events resulting in discontinuation were similar to those seen in previous studies with PEGETRON®. Only anemia and fatigue were reported as events that led to discontinuation in > 1% of patients in any arm.

Adverse reactions that led to dose modifications of any medication (primarily PEGETRON®) occurred in 39% of patients who received VICTRELIS TRIPLE™ compared to 24% of patients who received PEGETRON® alone. The most common reason for dose reduction was anemia, which occurred more frequently in patients who received VICTRELIS TRIPLE™ than in patients who received PEGETRON® alone.

Adverse reactions were considered by investigators to be causally related to VICTRELIS TRIPLE™. Adverse reactions reported in ≥ 4% of patients who received VICTRELIS TRIPLE™ and reported at a greater rate than the PEGETRON® in SPRINT-1, SPRINT-2, and RESPOND-2 are listed by System Organ Class in Table 2.

Table 2: Treatment-related Adverse Reactions Reported in $\geq 4\%$ of Patients who Received VICTRELIS TRIPLE™ and Reported at a Greater Rate than PEGETRON® alone^a

Adverse Reactions	Previously Untreated (SPRINT-1 & SPRINT-2)		Previous Treatment Failures (RESPOND-2)	
	Patients Reporting Adverse Reactions		Patients Reporting Adverse Reactions	
Body System Organ Class	BOC/PegIFN α 2b/ RBV n = 1,225 (%)	PegIFN α 2b/RBV n = 467 (%)	BOC/PegIFN α 2b/ RBV n = 323 (%)	PegIFN α 2b/RBV n = 80 (%)
Median Exposure (days)	197	216	253	104
Blood and Lymphatic System Disorders				
Anemia	50	30	45	20
Leukopenia	9	8	5	1
Neutropenia	25	19	14	10
Thrombocytopenia	4	1	3	0
Eye Disorders				
Vision Blurred	7	5	2	1
Gastrointestinal Disorders				
Abdominal Pain	5	4	3	9
Abdominal Pain Upper	7	7	6	3
Constipation	6	5	8	5
Diarrhea	23	19	23	15
Dry Mouth	10	9	14	9
Dysgeusia	35	16	44	11
Dyspepsia	7	7	6	5
Gastroesophageal Reflux Disease	5	2	5	0
Nausea	45	40	41	38
Stomatitis	4	3	4	3
Vomiting	19	12	13	8
General Disorders and Administration Site Conditions				
Asthenia	15	18	21	16
Chills	33	29	33	30
Fatigue	58	58	55	50
Pain	10	8	7	4
Pyrexia	32	32	28	21
Investigations				
Decreased Weight	11	12	11	9
Metabolism and Nutrition Disorders				
Decreased Appetite	25	24	25	16
Musculoskeletal and Connective Tissue Disorders				
Arthralgia	18	17	21	14
Back Pain	6	6	6	4
Muscle Spasms	3	3	4	3
Nervous System Disorders				
Dizziness	18	14	16	10
Headache	45	42	40	48
Memory Impairment	4	5	5	4
Paresthesia	4	2	3	1
Psychiatric Disorders				
Anxiety	13	12	12	6
Depression	23	22	16	15

Adverse Reactions	Previously Untreated (SPRINT-1 & SPRINT-2)		Previous Treatment Failures (RESPOND-2)	
	Patients Reporting Adverse Reactions		Patients Reporting Adverse Reactions	
Body System Organ Class	BOC/PegIFN α 2b/ RBV n = 1,225 (%)	PegIFN α 2b/RBV n = 467 (%)	BOC/PegIFN α 2b/ RBV n = 323 (%)	PegIFN α 2b/RBV n = 80 (%)
Insomnia	33	33	29	20
Irritability	22	23	21	13
Mood Altered	4	3	2	3
Respiratory, Thoracic, and Mediastinal Disorders				
Cough	18	22	23	16
Dyspnea	27	23	33	21
Epistaxis	3	2	5	3
Skin and Subcutaneous Tissue Disorders				
Alopecia	27	27	22	16
Dry Skin	17	18	22	8
Pruritus	24	25	21	18
Rash	18	20	16	5
Rash Maculo-papular	4	3	3	0
Rash Papular	5	3	1	0

BOC = boceprevir; PegIFN α 2b = peginterferon alfa-2b; RBV = ribavirin
a: Injection-site reactions have not been included since VICTRELIS™ is administered orally.

Serious Adverse Drug Reactions

Serious AEs were reported in 11% of patients who received VICTRELIS TRIPLE™ and 8% of patients who received PEGETRON®.

There were eight patient deaths that occurred during the treatment or follow-up in these clinical trials. Four deaths occurred in patients who received PEGETRON® alone (4/547, 1%), and 4 deaths occurred in the patients who received VICTRELIS TRIPLE™ (4/1,548, < 1%).

Anemia

Anemia was observed in 49% of patients treated with VICTRELIS TRIPLE™ compared with 29% of patients treated with PEGETRON® alone. With the interventions used for anemia management in the clinical trials, the average additional decrease of hemoglobin was approximately 10 g/L in hemoglobin concentration when VICTRELIS™ was added to PEGETRON®. The mean decreases in hemoglobin values from baseline were larger in previously treated patients compared to patients who had never received prior therapy.

Dose modifications (generally of PEGETRON®) due to anemia/hemolytic anemia occurred twice as often in patients treated with VICTRELIS TRIPLE™ compared to PEGETRON® alone (13%). The proportion of patients who received a transfusion for the management of anemia was 3% of patients in the VICTRELIS TRIPLE™ arms compared to < 1% of patients who received PEGETRON® alone. Although not approved for the treatment of anemia associated with Hepatitis C, the use of Erythropoiesis Stimulating Agents (ESAs) was permitted for management of anemia, at the investigator's discretion, with or without RBV dose reduction in the Phase 2 and 3 clinical trials.

The proportion of subjects who received an ESA was 43% in the VICTRELIS TRIPLE™ arms compared to 24% in the PEGETRON® arm.

PEGETRON®

General Safety Information

In clinical trials, approximately 1.2% of patients treated with PEGETRON® reported life-threatening psychiatric events during treatment. These events included suicide, attempted suicide, suicidal ideation, psychosis including hallucinations, bipolar disorders and aggressive behavior, sometimes directed towards others (see **WARNINGS AND PRECAUTIONS**).

As with other alpha interferons, ophthalmological disorders including retinopathies (including macular edema), retinal hemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilledema have been rarely reported during therapy with PegIFN α 2b (see **WARNINGS AND PRECAUTIONS**).

Adverse reactions of the cardiovascular system (CVS), particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior cardiotoxic therapy. Cardiomyopathy was also observed in patients treated with PegIFN α peginterferon alpha and has been reported more frequently in patients with known risk factors for cardiovascular diseases. There are limited data to assess the reversibility of cardiomyopathy reported with the use of PegIFN α peginterferon alpha; however cases of reversible cardiomyopathy have been reported with the use of interferon alpha.

As with other alpha interferons, seizures, pancreatitis, hypertriglyceridemia, arrhythmia, diabetes, peripheral neuropathy, colitis (including ischemic and ulcerative), aplastic anemia, hypertension, cardiac ischemia, myocardial infarction, cerebrovascular ischemia, cerebrovascular hemorrhage, encephalopathy (see **WARNINGS AND PRECAUTIONS**), sarcoidosis or exacerbation of sarcoidosis, erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis, rhabdomyolysis, myositis, renal failure and renal insufficiency have been rarely or very rarely reported during therapy with PegIFN α 2b.

Very rarely, alpha interferons, including PegIFN α 2b used alone or in combination with RBV, may be associated with Pure Red Cell Aplasia (PRCA).

Alpha interferons have been associated with altered lipid metabolism (including hypercholesterolemia and hyperlipemia) and pulmonary hypertension.

Adequate hydration must be maintained in patients undergoing PegIFN α 2b therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

Refer to the Product Monograph for PEGETRON® for further detailed information.

Clinical Trial Adverse Drug Reactions ($\geq 1\%$ to $< 4\%$)

Adverse reactions reported in $< 4\%$ and $\geq 1\%$ of patients who received VICTRELIS TRIPLE™

VICTRELIS TRIPLE™ (boceprevir, ribavirin plus peginterferon alfa-2b)

Page 20 of 97

reported at a greater rate than the PEGETRON[®] arms in SPRINT-1, SPRINT-2, and RESPOND-2 are listed by System Organ Class in Table 3.

Table 3: Adverse Drug Reactions reported in $\geq 1\%$ to $< 4\%$ of Previously Untreated Patients (SPRINT-1 and SPRINT-2) and Previous Treatment Failures (RESPOND-2); reported at rates greater than the PEGETRON[®] Arms

Body System Organ Class / Adverse Reactions	Previously Untreated (SPRINT-1 & SPRINT-2)	Previous Treatment Failures (RESPOND-2)
	BOC/PegIFN α 2b/RBV Patients Reporting Adverse Reactions (%)	BOC/PegIFN α 2b/RBV Patients Reporting Adverse Reactions (%)
Cardiac Disorders		
Tachycardia	1	1
Ear and Labyrinth Disorders		
Tinnitus	3	2
Endocrine Disorders		
Hypothyroidism	3	2
Eye Disorders		
Retinal Exudates	1	1
Vision Impairment	2	1
Gastrointestinal Disorders		
Abdominal Discomfort	3	2
Abdominal Distention	1	2
Anorectal Discomfort	1	1
Aphthous Stomatitis	3	2
Cheilitis	2	1
Flatulence	2	2
Gingivitis	< 1	2
Glossodynia	2	2
Hemorrhoids	2	1
Mouth Ulceration	2	2
Oral Pain	2	1
Tongue Ulceration	1	2
Tooth Disorder	1	3
General Disorders and Administration Site Conditions		
Chest Pain	2	1
Feeling of Body Temperature Change	< 1	1
Malaise	2	2
Mucosal Dryness	1	2
Infections and Infestations Disorders		
Cellulitis	1	1
Herpes Simplex	2	2
Influenza	1	1
Oral Fungal Infection	2	3
Sinusitis	2	1
Urinary Tract Infection	1	1
Metabolism and Nutrition Disorders		
Dehydration	1	1
Hypertriglyceridemia	1	1

Body System Organ Class / Adverse Reactions	Previously Untreated (SPRINT-1 & SPRINT-2)	Previous Treatment Failures (RESPOND-2)
	BOC/PegIFN α 2b/RBV Patients Reporting Adverse Reactions (%)	BOC/PegIFN α 2b/RBV Patients Reporting Adverse Reactions (%)
Musculoskeletal and Connective Tissue Disorders		
Neck Pain	1	2
Nervous System Disorders		
Amnesia	1	< 1
Hypoesthesia	2	1
Parosmia	1	3
Syncope	1	2
Psychiatric Disorders		
Affect lability	3	2
Aggression	1	1
Anger	1	2
Confusional State	1	< 1
Libido Disorder	2	2
Suicidal Ideation	1	1
Renal and Urinary Disorders		
Pollakiuria	2	1
Reproductive System and Breast Disorders		
Erectile dysfunction	1	1
Respiratory, Thoracic, and Mediastinal Disorders		
Dry Throat	1	1
Oropharyngeal Pain	3	2
Respiratory Tract Congestion	1	2
Sinus Congestion	1	3
Wheezing	1	1
Skin and Subcutaneous Tissue Disorders		
Dermatitis	2	3
Eczema	3	2
Edema Peripheral	2	2
Erythema	2	3
Rash Erythematous	3	1
Skin Lesion	1	1
Vascular Disorders		
Hypotension	1	2
BOC = boceprevir; PegIFN α 2b = peginterferon alfa-2b; RBV = ribavirin		

Less Common Clinical Trial Adverse Drug Reactions (< 1%)

Adverse reactions reported in < 1% of patients who received VICTRELIS TRIPLE™ reported at a greater rate than the PEGETRON® in SPRINT-1, SPRINT-2, and RESPOND-2 are listed.

Blood and Lymphatic System Disorders: Hemorrhagic diathesis, hemolysis, lymphadenopathy, and lymphopenia

Cardiac Disorders: Acute myocardial infarction, arrhythmia, atrial fibrillation, cardiovascular

disorder, coronary artery disease, palpitations, pericardial effusion, and pericarditis

Ear and Labyrinth Disorders: Deafness, ear discomfort, and hearing impaired

Endocrine Disorders: Goiter

Eye Disorders: Abnormal sensation in eye, conjunctival hemorrhage, conjunctivitis, eye pain, eye pruritus, eye swelling, eyelid edema, increased lacrimation, ocular hyperemia, papilledema photophobia, retinal ischemia, and retinopathy

Gastrointestinal Disorders: Anal pruritus, colitis, dry lip, dysphagia, feces discolored, frequent bowel movements, gastritis, gingival bleeding, gingival pain, glossitis, lower abdominal pain, odynophagia, pancreatic insufficiency, pancreatitis, proctalgia, rectal hemorrhage, salivary hypersecretion, sensitivity of teeth, and tongue discoloration

Hepatobiliary Disorders: Cholecystitis

General Disorders and Administration Site Conditions: Chest discomfort, impaired healing, and non-cardiac chest pain

Immune System Disorders: Sarcoidosis and non-acute porphyria

Infections and Infestations Disorders: Ear infection, epiglottitis, fungal skin infection, gastroenteritis, onychomycosis, otitis media, pharyngitis, respiratory tract infection, rhinitis, sepsis, and skin infection

Investigations: Cardiac murmur

Metabolism and Nutrition Disorders: Appetite disorder, diabetes mellitus, gout, hypercalcemia, and hypokalemia

Musculoskeletal and Connective Tissue Disorders: Arthritis, bone pain, joint swelling, muscular weakness, musculoskeletal chest pain, and musculoskeletal pain

Neoplasms Benign, Malignant and Unspecified (including cysts and polyps): Thyroid neoplasm

Nervous System Disorders: Cerebral ischemia, encephalopathy, hyperesthesia, mental impairment, neuralgia, neuropathy peripheral, and pre-syncope

Psychiatric Disorders: Abnormal behaviour, agitation, apathy, auditory hallucination, bipolar disorder, completed suicide, homicidal ideation, mental status changes, panic attack, paranoia, psychiatric decompensation, restlessness, suicide attempt, suicidal ideation, and visual hallucination

Renal and Urinary Disorders: Dysuria and nocturia

Reproductive System and Breast Disorders: Amenorrhea and aspermia

Respiratory, Thoracic and Mediastinal Disorders: Dysphonia, increased upper airway secretion, oropharyngeal blistering, orthopnea, pleural fibrosis, pleuritic pain, pulmonary embolism, and respiratory failure

Skin and Subcutaneous Tissue Disorders: Photosensitivity reaction, skin ulcer, and urticaria

Vascular Disorders: Blood pressure fluctuation, deep vein thrombosis, flushing, pallor, peripheral coldness, and venous thrombosis

Abnormal Hematologic and Clinical Chemistry Findings

Changes in selected laboratory values during treatment of adult patients with VICTRELIS TRIPLE™ are described in Table 4. Decreases in hemoglobin may require a decrease in dosage or interruption of RBV (see **WARNINGS AND PRECAUTIONS** and **CLINICAL TRIALS**).

Table 4: Selected Laboratory Values

Hematological Parameters	Previously Untreated (SPRINT-1 & SPRINT-2)		Previous Treatment Failures (RESPOND-2)	
	Patients Reporting Selected Hematological Parameters		Patients Reporting Selected Hematological Parameters	
	BOC/PegIFNα2b/RBV n = 1,225 (%)	PegIFNα2b/RBV n = 467 (%)	BOC/PegIFNα2b/RBV n = 323 (%)	PegIFNα2b/RBV n = 80 (%)
Hemoglobin (g/L)				
< 100	49	29	49	25
< 85	6	3	10	1
Platelets (x 10⁹/L)				
< 50	3	1	4	0
< 25	< 1	0	0	0
Neutrophils (x 10⁹/L)				
< 0.75	31	18	26	13
< 0.5	8	4	7	4

BOC = boceprevir; PegIFNα2b = peginterferon alfa-2b; RBV = ribavirin

An increase in uric acid and indirect bilirubin values associated with hemolysis was observed in some patients treated with PEGETRON[®], but values returned to baseline levels by four weeks after the end of therapy. Among those patients with elevated uric acid levels, very few patients treated with PEGETRON[®] developed clinical gout, none of which required treatment modification or discontinuation from the clinical trials for PEGETRON[®].

Neutrophils and Platelets

The proportion of patients with decreased neutrophil and platelet counts was higher in the VICTRELIS TRIPLE™ arms compared to patients who received only PEGETRON[®].

Seven percent of patients who received VICTRELIS TRIPLE™ had neutrophil counts of < 0.5 x 10⁹/L compared to 4% of patients who received only PEGETRON[®]. Three percent of patients

who received VICTRELIS TRIPLE™ had platelet counts of $< 50 \times 10^9/L$ compared to 1% of patients who received only PEGETRON®. Three percent of patients who received VICTRELIS TRIPLE™ had platelet counts of $< 50 \times 10^9/L$ compared to 1% of patients who received only PEGETRON® (see **WARNINGS AND PRECAUTIONS, Hematologic, Neutropenia and ADVERSE REACTIONS, Abnormal Hematologic and Clinical Chemistry Findings**).

Post-Market Adverse Drug Reactions

VICTRELIS™

The post-marketing ADRs are not yet available for VICTRELIS™.

PEGETRON®

The following post-market adverse drug reactions have been reported with PEGETRON® alone.

Very rarely, post-marketing reports of the following ADRs have been received:

Cardiac Disorders: cardiac ischemia, myocardial infarction

Ear and Labyrinth Disorders: hearing impairment, hearing loss, vertigo

Gastrointestinal Disorders: colitis (ulcerative and ischemic), pancreatitis, stomatitis

General Disorders and Administration Site Conditions: injection site necrosis

Musculoskeletal and Connective Tissue Disorders: myositis, rhabdomyolysis

Nervous System Disorders: cerebrovascular hemorrhage, cerebrovascular ischemia, seizures

Renal and Urinary Disorders: renal failure, renal insufficiency

Skin and Subcutaneous Tissue Disorders: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including the following ADRs:

Blood and Lymphatic System Disorders: idiopathic thrombocytopenic purpura, thrombotic thrombocytopenic purpura

Eye Disorders: Vogt-Koyanagi-Harada syndrome

Musculoskeletal and Connective Tissue Disorders: rheumatoid arthritis, systemic lupus erythematosus

Vascular Disorders: vasculitis

Very rare cases of the following ADR have been received:

Blood and Lymphatic System Disorders: Pure Red Cell Aplasia (PRCA) were reported in patients who received both anti-hepatitis C therapy and ESAs. Erythropoiesis Stimulating Agents are not approved for the management of anemia associated with hepatitis C. The potential for an increased risk of PRCA and development of anti-erythropoietin (EPO) antibodies should therefore be kept in mind when considering EPO stimulating agents use for hepatitis C treatment induced-anemia.

The following cases of acute hypersensitivity reactions have been reported:

Immune System Disorders: anaphylaxis

Skin and Subcutaneous Tissue Disorders: angioedema, urticaria

Other AEs reported with PegIFN α 2b alone or in combination with RBV include the following ADRs:

Endocrine Disorders: hyperthyroidism, hypothyroidism

Eye Disorders: serous retinal detachment

Gastrointestinal Disorders: abdominal pain

General Disorders and Administration Site Conditions: asthenic conditions (including asthenia, malaise and fatigue), chest pain, irritability

Infections and Infestations Disorders: bacterial infection (including sepsis), fungal infection

Metabolism and Nutrition Disorders: dehydration

Nervous System Disorders: facial palsy, migraine headache, paresthesia, peripheral neuropathy

Psychiatric Disorders: anxiety, emotional lability

Respiratory, Thoracic and Mediastinal Disorders: cough, dyspnea

Skin and Subcutaneous Tissue Disorders: dry skin, pruritus, rash

Vascular Disorders: hypertension, hypotension, palpitations

DRUG INTERACTIONS

Serious Drug Interactions

Contraindicated Drugs: alfuzosin, amiodarone, propafenone, quinidine, carbamazepine, phenobarbital, phenytoin, rifampin, dihydroergotamine, ergonovine, ergotamine, methylergonovine, cisapride, St. John's Wort, lovastatin, simvastatin, sildenafil or tadalafil when used for the treatment of pulmonary arterial hypertension, pimozide, drospirenone, astemizole, terfenadine, midazolam (orally administered), and triazolam (orally administered) (see **CONTRAINDICATIONS**).

Overview

VICTRELIS™

Effect of Other Drugs on VICTRELIS™ Pharmacokinetics

VICTRELIS™ is partly metabolized by CYP3A4/5. Co-administration of VICTRELIS™ with medicines that induce or inhibit CYP3A4/5 could increase or decrease exposure to VICTRELIS™.

Effects of VICTRELIS™ on Pharmacokinetics of Other Drugs

VICTRELIS™ is a strong inhibitor of CYP3A4/5. Medicines metabolized primarily by CYP3A4/5 may have increased exposure when administered with VICTRELIS™, which could increase or prolong their therapeutic and adverse effects (see Table 5). VICTRELIS™ does not inhibit CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 or CYP2E1 *in vitro*. In addition, VICTRELIS™ does not induce CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19 or CYP3A4/5 *in vitro*.

PEGETRON®

Ribavirin: Results of *in vitro* studies using both human and rat liver microsome preparations indicated no cytochrome P450 enzyme-mediated metabolism of RBV. Ribavirin does not inhibit

cytochrome P450 enzymes. There is no evidence from toxicity studies that RBV induces liver enzymes. Therefore there is minimal potential for P450 enzyme-based interactions.

Peginterferon alfa-2b: Results of a single dose study with PegIFN α 2b demonstrated no effect on the activity of cytochrome P450 isoenzymes CYP1A2, CYP2C8/9, CYP2D6, and CYP3A4 or hepatic N-acetyl transferase. The literature, however, reports up to a 50% reduction in clearance of CYP1A2 substrates (e.g., theophylline) when administered with other forms of interferon alpha and therefore caution should be exercised when PegIFN α 2b is used with medications metabolized by CYP1A2.

A multi-dose probe study assessing P450 substrates was performed in 26 patients with chronic hepatitis C, who received a once-weekly PegIFN α 2b (1.5 mcg/kg) for 4 weeks. There was no inhibition of CYP1A2, 3A4 or N-acetyltransferase. There was a 27% increase in activity of CYP2C8/9 and a 69% increase in CYP2D6. Caution should be used when administering PegIFN α 2b with medications metabolized by CYP2C8/9 and CYP2D6.

No PK interactions were noted between PegIFN α 2b and RBV in a multiple-dose PK study.

Drug-Drug Interactions

VICTRELIS™

VICTRELIS™ is contraindicated when co-administered with medicines that are potent inducers of CYP3A4/5 and medicines that are highly dependent on CYP3A4/5 for clearance, and for which elevated plasma concentrations are associated with serious and/or life-threatening events (see Table 1, **CONTRAINDICATIONS**).

Boceprevir is metabolized primarily by aldo-ketoreductase (AKR). In drug interaction trials conducted with AKR inhibitors diflunisal and ibuprofen, BOC exposure did not increase to clinically significant extent. VICTRELIS™ may be administered with AKR inhibitors.

Boceprevir is partly metabolized by CYP3A4/5. *In vitro* BOC has been shown to be also a substrate of p-glycoprotein. Co-administration of VICTRELIS™ with drugs that induce or inhibit CYP3A4/5 could decrease or increase exposure to BOC. Co-administration of VICTRELIS™ is contraindicated with medicines that are potent CYP3A4/5 inducers, where significantly reduced BOC plasma concentrations may be associated with reduced efficacy (see Table 1, **CONTRAINDICATIONS**).

Table 5 provides recommendations based on established, potential drug interactions or drugs contraindicated with VICTRELIS™.

Table 5: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction (Tables 24 and 25)

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
<i>Antiarrhythmics</i>		
bepridil	↑ bepridil	Co-administration of VICTRELIS™ with bepridil has the potential to produce serious and/or life-threatening AEs and has not been studied. Caution is warranted and therapeutic concentration monitoring of these drugs is recommended if they are used concomitantly with VICTRELIS™.
digoxin	↑ digoxin	Digoxin concentrations may be increased with VICTRELIS™. Use the lowest dose initially with careful titration and monitoring of serum digoxin concentrations.
<i>Anticoagulants</i>		
warfarin	↑ or ↓ warfarin	Concentrations of warfarin may be altered when co-administered with VICTRELIS™. It is recommended that international normalization ratio be monitored when warfarin is co-administered with VICTRELIS™.
<i>Antidepressants</i>		
desipramine trazodone	↑ desipramine ↑ trazodone	Plasma concentrations of desipramine and trazodone may increase when administered with VICTRELIS™, resulting in AEs such as dizziness, hypotension and syncope. Use with caution and consider a lower dose of desipramine or trazodone.

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
Antifungals		
<p style="text-align: center;"> ketoconazole itraconazole posaconazole voriconazole </p>	<p style="text-align: center;"> ↑ BOC ↑ ketoconazole ↑ itraconazole ↑ posaconazole ↑ voriconazole </p>	<p>The results of the drug interaction study between ketoconazole (400 mg twice daily (BID)) and BOC (400 mg single dose) led to increased plasma exposure of BOC (AUC, 131% ↑, C_{max} 41% ↑). These changes were not considered clinically significant and no dose adjustment for BOC or ketoconazole is required.</p> <p>Plasma concentrations of ketoconazole, itraconazole, voriconazole or posaconazole may be increased with VICTRELIS™. When co-administration is required, doses of ketoconazole and itraconazole should not exceed 200 mg/day.</p>
Anti-gout		
<p style="text-align: center;">colchicine</p>	<p style="text-align: center;">↑ colchicine</p>	<p>Significant increases in colchicine levels are expected; fatal colchicine toxicity has been reported with other strong CYP3A4 inhibitors.</p> <p>Patients with renal or hepatic impairment should not be given colchicine with VICTRELIS™.</p> <p>Treatment of gout flares (during treatment with VICTRELIS™): 0.6 mg (1 tablet) x 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Dose to be repeated no earlier than 3 days.</p> <p>Prophylaxis of gout flares (during treatment with VICTRELIS™): If the original regimen was 0.6 mg twice a day, reduce dose to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, reduce the dose to 0.3 mg once every other day.</p> <p>Treatment of familial Mediterranean fever (FMF) (during treatment with VICTRELIS™): Maximum daily dose of 0.6 mg (maybe given as 0.3 mg twice a day).</p>

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
<i>Anti-infective</i>		
clarithromycin	↑ clarithromycin	Concentrations of clarithromycin may be increased with VICTRELIS™; however, no dosage adjustment is necessary for patients with normal renal function.
<i>Antimycobacterials</i>		
rifabutin	↓ BOC ↑ rifabutin	Increases in rifabutin exposure are anticipated, while exposure of BOC may be decreased. Doses have not been established when rifabutin is used in combination with VICTRELIS™. Concomitant use rifabutin with VICTRELIS™ is not recommended.
<i>Calcium Channel Blockers, dihydropyridine</i>		
felodipine nifedipine nicardipine	↑ dihydropyridine calcium channel blockers	Plasma concentrations of dihydropyridine calcium channel blockers may increase when administered with VICTRELIS™. Caution is warranted and clinical monitoring is recommended.
<i>Corticosteroid, inhaled</i>		
budesonide fluticasone	↑ budesonide ↑ fluticasone	Concomitant use of inhaled budesonide or fluticasone with VICTRELIS™ may result in increased plasma concentrations of budesonide or fluticasone, resulting in significantly reduced serum cortisol concentrations. Avoid co-administration if possible, particularly for extended durations.
<i>Corticosteroid, systemic</i>		
dexamethasone	↓ BOC	Co-administration of VICTRELIS™ with CYP3A4/5 inducers may decrease plasma concentrations of BOC, which may result in loss of therapeutic effect. Therefore, this combination should be avoided if possible and used with caution if necessary.
<i>Endothelin Receptor Antagonist</i>		
bosentan	↑ bosentan	Concentrations of bosentan may be increased when co-administered with VICTRELIS™. Use with caution and monitor closely.

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
<i>HCV Antivirals</i>		
PegIFNα2b	<p style="text-align: center;">↔ BOC ↔ PegIFNα2b</p>	<p>The results of the drug interaction study between PegIFNα2b (1.5 mcg/kg SC weekly) and BOC (400 mg TID) demonstrated that exposure to BOC and PegIFNα2b was not significantly affected when co-administered. No dose adjustment required for VICTRELIS™ or PegIFNα2b. The interaction between BOC and PEGIFNα2a has not been studied.</p>
<i>HIV-Antiviral Agents: Non-nucleoside Reverse Transcriptase Inhibitors</i>		
efavirenz	<p style="text-align: center;">↓ BOC ↔ efavirenz</p>	<p>The results of the drug interaction study between efavirenz (600 mg daily) and BOC (800 mg TID) demonstrated a decreased plasma trough concentrations of BOC (C_{\min} 44% ↓). The clinical outcome of this observed reduction of BOC trough concentrations has not been directly assessed.</p>
<i>HIV-Antiviral Agents: Nucleoside Reverse Transcriptase Inhibitors</i>		
tenofovir	<p style="text-align: center;">↔ BOC ↑ tenofovir</p>	<p>The results of the drug interaction study between tenofovir (300 mg daily) and BOC (800 mg TID) led to increased plasma exposure of tenofovir (C_{\max} 32% ↑). These changes were not considered clinically significant and no dose adjustment for VICTRELIS™ or tenofovir is required.</p>
<i>HIV-Antiviral Agents: HIV PIs</i>		
ritonavir	<p style="text-align: center;">↓ BOC ↑ or ↓ HIV PIs</p>	<p>The results of the drug interaction study between ritonavir (100 mg daily) and BOC (400 mg TID) demonstrated no notable change in the exposure of BOC (C_{\max} 27% ↓). These changes were not considered clinically significant and no dose adjustment for VICTRELIS™ or ritonavir is required. The effect of ritonavir-boosted HIV PIs on BOC exposure is unknown. The effect of BOC on HIV PI concentrations is unknown.</p>

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
<i>HMG-CoA Reductase Inhibitors</i>		
atorvastatin rosuvastatin pravastatin fluvastatin	↑ atorvastatin ↑ rosuvastatin ↑ pravastatin ↑ fluvastatin	Increases in atorvastatin and rosuvastatin exposures are anticipated. Titrate atorvastatin dose carefully and do not exceed maximum daily dose of 20 mg during co-administration with VICTRELIS™. For rosuvastatin, pravastatin and fluvastatin, the drug interaction effect has not been studied and, therefore, caution should be used.
<i>Immunosuppressants</i>		
cyclosporine sirolimus tacrolimus	↑ immunosuppressants	Plasma concentrations of cyclosporine, sirolimus and tacrolimus are expected to be increased significantly during co-administration with VICTRELIS™. Close monitoring of immunosuppressant blood levels is recommended.
<i>Inhaled beta-agonist</i>		
salmeterol	↑ salmeterol	Concentrations of salmeterol may be increased when co-administered with VICTRELIS™. Concurrent use of inhaled salmeterol and VICTRELIS™ is not recommended due to the risk of cardiovascular events associated with salmeterol, including QT prolongation, palpitations and sinus tachycardia.
<i>Narcotic Analgesic/Opioid Dependence</i>		
buprenorphine methadone	↑ or ↓ buprenorphine ↑ or ↓ methadone	Plasma concentrations of buprenorphine or methadone may increase or decrease when co-administered with VICTRELIS™. However, the combination has not been studied. Clinical monitoring is recommended as the dose of buprenorphine or methadone may need to be altered during concomitant treatment with VICTRELIS™.

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
<i>Non-Steroidal Anti-Inflammatories (NSAIDs)</i>		
<p style="text-align: center;">diflunisal</p> <p style="text-align: center;">ibuprofen</p>	<p style="text-align: center;">↔ BOC</p> <p style="text-align: center;">↔ BOC</p>	<p>The results of the drug interaction study between diflunisal (250 mg BID) and BOC (800 mg TID) demonstrated no substantive change in the exposure of BOC (C_{min} 31% ↑). These changes were not considered clinically significant and no dose adjustment for VICTRELIS™ or diflunisal is required.</p> <p>The results of the drug interaction study between ibuprofen (600 mg TID) and BOC (400 mg single dose) demonstrated no clinically relevant change. No dose adjustment for VICTRELIS™ or ibuprofen is required.</p>
<i>Oral Hormonal Contraceptives</i>		
<p style="text-align: center;">drospirenone/ethinyl estradiol</p>	<p style="text-align: center;">↑ drospirenone ↓ ethinyl estradiol</p>	<p>The results of the drug interaction study between BOC (800 mg TID) and oral drospirenone/ethinyl estradiol (3 mg/0.02 mg daily) at steady-state demonstrated an increased systemic exposure of drospirenone (AUC, 99%; C_{max}, 57%) without notably affecting the exposures of ethinyl estradiol (AUC, 24% ↓ and C_{max}, ↔). Therefore, alternative methods of non-hormonal contraception are recommended. Co-administration of VICTRELIS™ with drospirenone is contraindicated (see CONTRAINDICATIONS).</p>

Concomitant Drug Class/Name	Effect on Concentration of BOC or Concomitant Drug	Clinical Comment
<i>PDE-5 Enzyme Inhibitors</i>		
<p>sildenafil tadalafil</p>	<p>↑ sildenafil ↑ tadalafil</p>	<p>Increases in PDE-5 inhibitor concentrations are expected, and may result in an increase in AEs, including hypotension, syncope, visual disturbances, and priapism.</p> <p>Use of sildenafil or tadalafil when used for the treatment of PAH is contraindicated with VICTRELIS™ (see CONTRAINDICATIONS).</p> <p><u>Use of PDE-5 inhibitors for erectile dysfunction:</u></p> <p>Use with caution in combination with VICTRELIS™ with increased monitoring for PDE-5 inhibitor-associated AEs. Do not exceed the following doses: sildenafil: 25 mg every 48 hours tadalafil: 10 mg every 72 hours vardenafil: 2.5 mg every 24 hours (Dose not approved in Canada. Therefore, co-administration of vardenafil with VICTRELIS TRIPLE™ is not recommended.)</p>
<p>vardenafil</p>	<p>↑ vardenafil</p>	
<i>Sedatives/Hypnotics</i>		
<p>alprazolam (intravenous administration) midazolam (intravenous administration) triazolam (intravenous administration)</p>	<p>↑ alprazolam ↑ midazolam ↑ triazolam</p>	<p>No interaction studies have been done with intravenous benzodiazepines. Close clinical monitoring for respiratory depression and/or prolonged sedation should be exercised during co-administration of VICTRELIS™ with intravenous benzodiazepines (alprazolam, midazolam, triazolam). Dose adjustment of the benzodiazepine should be considered.</p>

BOC = boceprevir; PegIFN α 2b = peginterferon alfa-2b

PEGETRON®

Ribavirin was shown *in vitro* to inhibit phosphorylation of the nucleoside analog reverse transcriptase inhibitors (NRTIs) zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of RBV with either zidovudine or stavudine might lead to increased HIV plasma viremia. Therefore, it is recommended that plasma HIV-RNA levels be closely monitored in patients treated with RBV concurrently with either of these two agents. If HIV-RNA levels increase, the use of RBV concomitantly with NRTIs must be reviewed.

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, RBV increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogues (e.g., didanosine or abacavir). Co-administration of RBV and didanosine is not recommended. Reports of mitochondrial toxicity in particular lactic acidosis and pancreatitis, of which some fatal, have been reported.

An increased risk of developing peripheral neuropathy cannot be ruled out for treatments combining telbivudine with any alpha (standard or pegylated types) and beta interferon products. This risk might occur when the drug is used with interferon products other than pegylated interferon alfa-2a. Development of peripheral neuropathy was reported in a small clinical trial in Hepatitis B investigating the use of both 600 mg daily of telbivudine and 180 mcg once weekly by subcutaneous administration of pegylated interferon alfa-2a. The safety and efficacy of telbivudine in combination with interferon alfa-2b has not been established in patients with Hepatitis B; therefore, telbivudine in combination with alpha interferons is not recommended. There are limited data regarding the efficacy and safety of combining PEGETRON[®] with Highly Active Anti-Retroviral Therapy (HAART) for HIV-HCV co-infected patients. Patients co-infected with HIV and who are receiving HAART may be at increased risk of developing lactic acidosis. Caution should be used when treating HIV-HCV co-infected patients with PEGETRON[®] in combination with HAART.

Any potential for interactions may persist for up to two months (5 half-lives for RBV) after cessation of PEGETRON[®] therapy due to the long half-life of RBV.

There is no evidence that RBV interacts with non-nucleoside reverse transcriptase inhibitors (NNRTIs) or PIs.

Antacid: The bioavailability of RBV (600 mg) was decreased by co-administration with an antacid containing magnesium aluminum and simethicone; AUC_{0-∞} decreased 14%. This interaction is not considered to be clinically relevant.

Drug-Food Interactions

Increased exposure to BOC was observed following administration with food. VICTRELIS TRIPLE[™] should be taken with food. The type of food does not affect exposure to BOC (see **ACTION AND CLINICAL PHARMACOLOGY**).

Drug-Herb Interactions

As with other alpha interferons, pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in PegIFN α 2b treated patients. These symptoms have been reported more frequently when *Shosaikoto* (also known as *Xiao-Chai-Hu-Tang*), a Chinese herbal medication, has been administered concomitantly with alpha interferons.

St. John's wort (*Hypericum perforatum*)

Co-administration of VICTRELIS TRIPLE™ and St. John's wort may lead to loss of virologic response to VICTRELIS TRIPLE™ (see **CONTRAINDICATIONS**).

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Ribavirin Capsules are for oral administration only.

Peginterferon alfa-2b Powder for Solution is administered subcutaneously using the REDIPEN® single dose delivery system with the disposable needles supplied.

VICTRELIS™ (boceprevir) Capsules are for oral administration only. **It is important that the dose of VICTRELIS™ (800 mg) be taken three times a day (every 7–9 hours).**

VICTRELIS TRIPLE™ treatment should be initiated only by a physician experienced in the treatment of patients with hepatitis C.

Recommended Dose

When treating adult patients with genotype 1 HCV, the three treatment medicines use the following recommended dosing:

PEGETRON®

The recommended dose of RBV Capsules to be used is 800–1,400 mg daily based upon patient weight (see Table 6). Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

The recommended dose of PegIFNα2b Powder for Solution is 1.5 mcg/kg/week subcutaneously (see Table 6). Peginterferon alfa-2b should be administered once a week.

Table 6: PEGETRON® Dosing Recommendations for Patients who are previously untreated and who have failed previous therapy^a

	Patient Weight (kg)	PegIFNα2b Powder for Solution		Ribavirin Capsules	
		Weekly Dose (mcg/kg)	REDIPEN® Size (mcg/0.5 mL) ^b	Daily Dose (mg)	Number of Capsules (200 mg)
Previously Untreated Patients	40 to 50	1.5	80	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
	51 to 65	1.5	100	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
	66 to 80	1.5	120	1,000	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
	81 to 105	1.5	150	1,200	3 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
	> 105	1.5	*	1,400	3 x 200 mg capsules A.M. 4 x 200 mg capsules P.M.
Previous Treatment Failure	40 to 50	1.5	80	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
	51 to 65	1.5	100	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
	66 to 85	1.5	120	1,000	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
	86 to 105	1.5	150	1,200	3 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
	> 105	1.5	*	1,400	3 x 200 mg capsules A.M. 4 x 200 mg capsules P.M.

PegIFNα2b = peginterferon alfa-2b

a: The daily dose for ribavirin Capsules approximately falls within 13 ± 2 mg/kg/day.

b: When reconstituted as instructed

*: Should be calculated based on the body weight of an individual patient

VICTRELIS™

The recommended dose of VICTRELIS™ is 800 mg (four 200 mg capsules) administered orally TID (every 7–9 hours) with food (a meal or light snack).

Duration of Therapy

The recommendations for duration of therapy differ for some subgroups from the dosing durations studied in the Phase 3 clinical trials (see **CLINICAL TRIALS**). Response-Guided Therapy is recommended for most individuals, but longer dosing is recommended in target groups (e.g., patients with cirrhosis).

Discontinuation of Dosing Based on Treatment Futility

Discontinuation of therapy is recommended in all patients with:

- 1) HCV-RNA levels ≥ 100 IU/mL at TW 12; or
- 2) Confirmed detectable HCV-RNA levels at TW 24.

The following recommendations for duration of therapy are described as follows and stated in the subsequent paragraphs:

- Previously Untreated Patients
 - Patients without cirrhosis
 - Patients with poor interferon response
 - Patients with cirrhosis
- Patients who have failed previous therapy with peginterferon and RBV
 - Patients (partial responders and relapsers) without cirrhosis
 - Patients with previous null response
 - Patients with cirrhosis

Recommended Duration of Therapy for Previously Untreated Patients

Patients without Cirrhosis

Therapy with PEGETRON[®] should be initiated for 4 weeks (TWs 1-4) based on dosing recommendations listed in Table 6.

Therapy with VICTRELIS[™] should be added to the PEGETRON[®] regimen at TW 5. Based on the patient’s HCV-RNA levels at TW 8, TW 12, and TW 24, use the following Response-Guided Therapy (RGT) guidelines to determine duration of treatment (see Table 7).

Table 7: Duration of Therapy with VICTRELIS TRIPLE[™] using RGT Guidelines in Patients without Cirrhosis who are Previously Untreated

ASSESSMENT ^a (HCV-RNA Results ^b)		ACTION
At TW 8	At TW 24	
Undetectable	Undetectable	Stop VICTRELIS TRIPLE [™] at TW 28. Treatment is completed.
Detectable	Undetectable	1. Continue VICTRELIS TRIPLE [™] until TW 28, and then 2. Administer PEGETRON [®] until TW 48.

- a: **Treatment Futility: If the patient has HCV-RNA results ≥ 100 IU/mL at TW 12 or if the patient has confirmed detectable HCV-RNA at TW 24, then discontinue the three-medicine regimen.**
- b: In clinical trials, plasma HCV-RNA was measured using a Roche COBAS* TaqMan* assay with a lower limit of detection of 9.3 IU/mL and a lower limit of quantification of 25 IU/mL (see **WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests**).

Patients with Poor Interferon Response

Consideration should be given to treating previously untreated patients who are poorly interferon responsive (less than a 1.0- \log_{10} decline in HCV-RNA at TW 4 with PegIFN α /RBV alone) with 4 weeks of PEGETRON[®] using the dosing recommendations listed in Table 6 followed by 44 weeks of VICTRELIS[™] in combination with PEGETRON[®] (see **CLINICAL TRIALS**).

Patients with Cirrhosis

Patients with compensated cirrhosis should receive 4 weeks of PEGETRON[®] using dosing recommendations listed in Table 6 followed by 44 weeks VICTRELIS[™] in combination with PEGETRON.

Recommended Duration of Therapy for Previous Treatment Failure

Patients without Cirrhosis (Partial Responders or Relapsers)

Therapy with PEGETRON[®] should be initiated for 4 weeks (TWs 1-4) based on dosing recommendations listed in Table 6.

Therapy with VICTRELIS[™] should be added to the PEGETRON[®] regimen at TW 5. Based on the patient's HCV-RNA levels at TW 8, TW 12, and TW 24 use the following Response-Guided Therapy (RGT) guidelines to determine duration of treatment (see Table 8).

Table 8: Duration of Therapy using RGT Guidelines in Patients without Cirrhosis who are Previous Treatment Failures (previous partial responders and relapsers)^a

ASSESSMENT ^b (HCV-RNA Results ^c)		ACTION
At TW 8	At TW 24	
Undetectable	Undetectable	Stop VICTRELIS TRIPLE [™] at TW 36. Treatment is completed.
Detectable	Undetectable	1. Continue all three medications until TW 36, and then 2. Administer PEGETRON [®] until TW 48.

- a: Previous Treatment Failures to PegIFN α /RBV Therapy: Previous Partial responders (Patients with a decrease in HCV-RNA viral load $\geq 2\text{-log}_{10}$ by Week 12 but never achieved SVR); Relapsers (Patients with undetectable HCV-RNA at end of prior treatment with a subsequent detectable HCV-RNA in plasma).
- b: **Treatment Futility: If the patient has HCV-RNA results ≥ 100 IU/mL at TW 12 or if the patient has confirmed detectable HCV-RNA at TW 24, then discontinue the three-medicine regimen.**
- c: In clinical trials, plasma HCV-RNA was measured using a Roche COBAS* TaqMan* assay with a lower limit of detection of 9.3 IU/mL and a lower limit of quantification of 25 IU/mL (see **WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests**).

Patients with Previous Null Response

Response-Guided Therapy was not studied in subjects who had less than a 2-log_{10} HCV-RNA decline by TW 12 during prior therapy with PegIFN α /RBV. If considered for treatment, these subjects should receive 4 weeks of PEGETRON[®] using dosing recommendations listed in Table 6 followed by 44 weeks of VICTRELIS[™] in combination with PEGETRON[®] (see **CLINICAL TRIALS**).

Response-Guided Therapy is not recommended for patients with prior null response.

Patients with Cirrhosis

Patients with compensated cirrhosis should receive 4 weeks of PEGETRON[®] using dosing recommendations listed in Table 6 followed by 44 weeks VICTRELIS[™] in combination with PEGETRON.

Dose Modifications

PEGETRON[®]

In general, the PEGETRON[®] dosage may be adjusted according to the patient's tolerance to the medication. If severe adverse reactions or laboratory abnormalities develop during the course of treatment, the dosage of PegIFN α 2b and/or RBV should be modified or therapy should be temporarily discontinued until the adverse reactions abate.

If persistent or recurrent intolerance develops despite adequate dosage adjustment, or if the disease progresses rapidly, treatment with VICTRELIS TRIPLE™ should be discontinued.

Dose Modification for ribavirin Treatment –Hemoglobin Levels

In patients who are previously untreated or who are previous treatment failures, it is recommended that a patient whose hemoglobin level falls below 100 g/L have his/her RBV dose reduced (see Table 9). A patient whose hemoglobin level falls below 85 g/L should be permanently discontinued from VICTRELIS TRIPLE™ therapy.

Table 9: Ribavirin Capsules Dose Reduction

	Starting dose (mg)	Dose after 1 st dose reduction (mg)	Dose after 2 nd dose reduction, if required (mg)
Previously Untreated Patients	800	600	400
	1,000	800	600
	1,200	1,000	800
	1,400	1,000	800
Previous Treatment Failure	800	600	400
	1,000	800	600
	1,200	800	600
	1,400	1,000	800

For patients with a history of stable cardiovascular disease, a RBV dose reduction is required if the hemoglobin decreases by ≥ 20 g/L during any 4 week period. In addition, for these cardiac history patients, if the hemoglobin remains < 120 g/L after 4 weeks on a reduced dose, the patient should discontinue VICTRELIS TRIPLE™ therapy.

Dose Modification for peginterferon alfa-2b Treatment - Neutrophil Count

It is recommended that a patient whose neutrophil count falls below $0.75 \times 10^9/L$, have his/her PegIFN α 2b dose reduced (see Table 10). A patient whose neutrophil count falls below $0.5 \times 10^9/L$ should be permanently discontinued from VICTRELIS TRIPLE™ therapy.

Table 10: Peginterferon alfa-2b Powder for Solution Dose Reduction

	Starting dose (mcg/kg/week)	Dose after 1 st dose reduction (mcg/kg/week)	Dose after 2 nd dose reduction, if required (mcg/kg/week)
Previously Untreated Patients And Previous Treatment Failures	1.5	1	0.5

Dose Modification for PEGETRON[®] Treatment - Laboratory values

The following guidelines for dose modification were developed in clinical trials based on laboratory values:

Table 11: Dose Modification for Patient

		Reduce only ribavirin dose, if ^a :	Reduce only PegIFNα2b if ^b :	Discontinue combination therapy if:
Previously Untreated Patients And Previous Treatment Failures	Hemoglobin	< 100 g/L	-	< 85 g/L
	Hemoglobin in patients with history of stable cardiac disease	≥ 20 g/L decrease in hemoglobin during any 4-week period during treatment (permanent dose reduction)		< 120 g/L after 4 weeks of dose reduction
	White blood cell count	-	< 1.5 x 10 ⁹ /L	< 1.0 x 10 ⁹ /L
	Neutrophil count	-	< 0.75 x 10 ⁹ /L	< 0.5 x 10 ⁹ /L
	Platelet count	-	< 50 x 10 ⁹ /L	< 25 x 10 ⁹ /L
	Bilirubin – Direct	-	-	2.5 x upper limit of normal
	Bilirubin – Indirect	> 5 mg/dL	-	> 4 mg/dL (for > 4 weeks)
	Creatinine	-	-	> 2.0 mg/dL
ALT/AST	-	-	2 x baseline AND > 10 x upper limit of normal	

PegIFNα2b = peginterferon alfa-2b

a: Ribavirin dose reduction: 1st dose reduction of ribavirin for adult patients is by 200 mg/day except in patients who received the 1,400 mg; it is by 400 mg/day. If needed, 2nd dose reduction of ribavirin is by an additional 200 mg/day.

b: Peginterferon alfa-2b dose reduction: dose reduction of peginterferon alfa-2b should be implemented in increments of 0.5 mcg/kg/week.

VICTRELIS[™]

Dose reduction of VICTRELIS[™] is not recommended. VICTRELIS[™] must not be administered in the absence of PegIFNα and RBV.

Dose Adjustments in Special Populations

Renal Impairment

No dose adjustment of VICTRELIS[™] is required in patients with any degree of renal impairment (see **DETAILED PHARMACOLOGY**). Patients with severe renal dysfunction (creatinine clearance < 50 mL/min) must not be treated with VICTRELIS TRIPLE[™] (see **CONTRAINDICATIONS**).

Hepatic Impairment

No dose adjustment of VICTRELIS[™] is required for patients with mild, moderate or severe hepatic impairment (see **DETAILED PHARMACOLOGY**, **Pharmacokinetics**, **Special Populations and Conditions**, Hepatic Insufficiency). Safety and efficacy of VICTRELIS[™] have

not been studied in patients with decompensated cirrhosis. VICTRELIS TRIPLE™ is contraindicated in patients with hepatic decompensation (Child-Pugh score > 6 [class B and C]).

Missed Dose

PEGETRON®

Ribavirin

If a patient misses a dose of RBV, the missed dose should be taken as soon as possible during the same day. If an entire day has gone by, the patient should check with their doctor about what to do. The patient should not double the next dose.

Peginterferon alfa-2b

If a patient misses a dose of PegIFN α 2b, the missed dose should be taken as soon as possible during the same day or on the next day, and continue the dosing schedule provided to you by your doctor. If several days go by, the patient should check with their doctor about what to do. The patient should not double the next dose.

VICTRELIS™

If a patient misses a dose and it is less than 2 hours before the next dose is due, the missed dose should be skipped.

If patient misses a dose and it is 2 or more hours before the next dose is due, the patient should take the missed dose with food and resume the normal dosing schedule.

Concomitant Therapy

Acetaminophen has been used successfully to alleviate the symptoms of fever and headache, which can occur with PegIFN α 2b. The recommended acetaminophen dosage is 500 mg to 1 g given 30 minutes before administration of PegIFN α 2b. The maximum dosage of acetaminophen to be given is 1 g four times daily. In order to properly assess the source of fever, adjunctive acetaminophen should be limited to a maximum of 5 consecutive days unless otherwise specified by the prescribing physician.

Reconstitution of peginterferon alfa-2b

Before you inject PegIFN α 2b using the REDIPEN® single dose delivery system, the two cartridges must be activated to mix (reconstitute) the powder with the sterile diluent to form a solution. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the REDIPEN® click. The powder must be completely dissolved. The appropriate PegIFN α 2b dose should be properly dialed and injected subcutaneously.

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. Do not use if discoloration is present.

Peginterferon alfa-2b should be administered at room temperature.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

VICTRELIS™

Daily doses of 3,600 mg have been taken by healthy volunteers for 5 days without untoward symptomatic effects.

There is no specific antidote for overdose with VICTRELIS™. Treatment of overdose with VICTRELIS™ should consist of general supportive measures, including monitoring of vital signs, and observation of the patient's clinical status. VICTRELIS™ is not eliminated by dialysis.

PEGETRON®

There is limited experience with overdosage. The primary effects of overdose were an increased incidence and severity of AEs reported at the therapeutic doses of PEGETRON®. Serious adverse events reported in cases of PEGETRON® overdose include affect lability, anemia, anxiety, ataxia, bursitis, dehydration, depression, fatigue, hallucination, hallucination auditory, hyperesthesia, hypothyroidism, mental disorder, myalgia, nausea, non-accidental overdose, neutropenia, paranoia, pneumonia, pyrexia, suicidal ideation, suicide attempt, thyroid disorder, urinary tract infection, and vomiting. In cases of overdoses, symptomatic treatment and close observation of the patient are recommended (see **DOSAGE AND ADMINISTRATION, Dose modification**).

In RBV plus interferon alfa-2b clinical trials, the maximum overdose reported was 10g of RBV capsules (50 x 200 mg capsules) taken with a dose of 39 million units of interferon alfa-2b (13 subcutaneous injections of 3 million IU each). The patient was observed for two days in the emergency room during which time no AE from the overdose was noted.

Ribavirin: Absorption of RBV is generally complete after one hour. Treatment of overdose with RBV consists of general supportive measures including monitoring of vital signs and observation of clinical status of the patient. There is no specific antidote for overdose with RBV. Although no data currently exists, administration of activated charcoal may be used to aid in the removal of unabsorbed drug. Ribavirin concentration is essentially unchanged by hemodialysis.

Peginterferon alfa-2b: Distinction between the therapeutic dose of PegIFN α 2b and overdose has not been clearly defined. Symptoms of overdose may include amplification of the adverse effects, notably "flu-like" symptoms, leukopenia or thrombocytopenia and increased serum liver enzyme levels. The severity of the adverse reactions can be ameliorated by adjusting the dose level and schedule, or in some cases termination of PegIFN α 2b therapy. Cardiovascular side effects such as hypotension and arrhythmia may require supportive therapy.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

VICTRELIS™

Boceprevir: VICTRELIS™ is an inhibitor of the HCV NS3/4A protease. VICTRELIS™ covalently, yet reversibly, binds to the NS3/4A protease active site serine (Ser139) through a (alpha)-ketoamide functional group to inhibit viral replication in HCV-infected host cells.

PEGETRON®

The mechanism of inhibition of HCV-RNA by PEGETRON® therapy has not been established.

Ribavirin: The mechanism by which RBV exerts its effects against HCV is unknown.

At physiologic concentrations, neither RBV nor its intracellular nucleotide metabolites have been shown to inhibit HCV-specific enzymes or HCV replication. Oral formulations of RBV monotherapy have been investigated as therapy for chronic hepatitis C in several clinical studies showing that RBV monotherapy had no effect on eliminating serum HCV or improving hepatic histology after 6 to 12 months of therapy and 6 months of follow up. However, when used in combination with PegIFN α 2b in the treatment of chronic hepatitis C, RBV has been shown to increase the efficacy of PegIFN α 2b used alone, as measured by reduction of viral load.

Peginterferon alfa-2b: Peginterferon alfa-2b is a covalent conjugate of recombinant interferon alfa-2b with monomethoxy polyethylene glycol (PEG, with an average molecular weight of 12,000 daltons). The average molecular weight of the conjugated molecule is approximately 31,000 daltons. *In vitro* and *in vivo* studies suggest that the biological activity of PegIFN α 2b is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type I interferons. The results of several studies suggest that, once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulation activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Pharmacodynamics

VICTRELIS™

Evaluation of Effect of VICTRELIS™ on QTc Interval

In a randomized, multiple-dose, placebo and active controlled four-way cross-over study, BOC was evaluated for the effect on QT/QTc intervals at the 800 mg TID (therapeutic dose) and the 1,200 mg TID dose in 36 healthy subjects after multiple dosing for 5 days. The mean maximum concentrations at the 800 and 1,200 mg doses were 1,690 ng/mL and 1,940 ng/mL, respectively.

There was no significant difference in the QTc interval between BOC and placebo. At the mean maximum BOC exposures of 1,690 and 1,940 ng/mL, which were achieved at 2 hours postdose for the 800 and 1,200 mg doses, the mean increases in placebo-adjusted QTcF were 4.5 and 0.3 ms with the upper limits of the 95% CI of 7.3 and 3.1 ms, respectively. The maximum observed mean increases in placebo-adjusted QTcF occurred at 4 hours post-dose, and were 5.8 and 2.9 ms with the upper limits of the 95% CI of 8.7 and 5.7 ms, respectively. Therefore, in this study with demonstrated ability to detect small effects, the upper bound of the one-sided 95% confidence interval for the largest placebo-adjusted, baseline-corrected QTc based on individual correction method was below 10 ms, the threshold for regulatory concern. The dose of 1,200 mg yields a BOC maximum exposure increase of approximately 15% which may not cover exposures due to co-administration with strong CYP3A4 inhibitors or use in patients with severe hepatic impairment. However, at the doses studied in the thorough QT study, no apparent concentration-QT relationship was identified. Thus, there is no expectation of a QTc effect under a higher exposure scenario.

Pharmacokinetics

Boceprevir

VICTRELIS™ capsules contain an approximately equal mixture of two diastereoisomers. In the plasma the diastereoisomer ratio is about 2:1 in favour of the active diastereoisomer, SCH 534128. The plasma concentrations of BOC described below consist of both diastereoisomers.

The PK properties of BOC have been evaluated in healthy adult subjects and HCV-infected patients (see Table 12 below and Table 22, **DETAILED PHARMACOLOGY, Pharmacokinetics**).

Table 12: Summary of boceprevir's PK Parameters at Steady-State in Healthy Subjects (n = 71)

Dose (mg)	C_{max} (ng/mL)	t_½ (h)	AUC_(τ) (ng·h/mL)	Clearance (L/h)	Volume of Distribution (L)
800 TID	1,723	3.0	5,408	159	717

AUC_(τ) = area under the plasma concentration-time curve from time 0 dosing interval

The PK profiles of single and multiple doses of BOC from 50 mg up to 800 mg, and 100 mg up to 1,200 mg, respectively, have been evaluated. In general, the PK results were similar between healthy and HCV subjects.

Absorption:

Boceprevir was absorbed following oral administration with a median T_{max} of 2 hours. Steady state AUC, C_{max} and C_{min} increased in a less-than dose-proportional manner and individual exposures overlapped substantially at 800 mg and 1,200 mg, suggesting diminished absorption at higher doses. Accumulation is minimal and PK steady state is achieved after approximately 1 day of TID dosing.

The absolute bioavailability of VICTRELIS™ has not been studied.

Effects of Food on Oral Absorption

VICTRELIS™ should be administered with food. Food enhanced the exposure of BOC by up to 60% at the 800 mg TID dose when administered with a meal, relative to the fasting state. The bioavailability of BOC was similar regardless of meal type (e.g., high-fat vs. low-fat) or whether taken 5 minutes prior to eating, during a meal, or immediately following completion of the meal. Therefore, VICTRELIS™ may be taken without regard to either meal type or timing.

Distribution:

Boceprevir has a mean apparent volume of distribution (Vd/F) of approximately 717 L (n = 71) at steady state. Human plasma protein binding is approximately 75% following a single dose of VICTRELIS™ 800 mg. Boceprevir is administered as an approximately equal mixture of two diastereomers which rapidly interconvert in plasma: one diastereomer is pharmacologically active and the other diastereomer is inactive.

Metabolism:

Studies *in vitro* indicate that BOC primarily undergoes metabolism through the AKR-mediated pathway to ketone-reduced metabolites that are inactive against HCV. After a single 800 mg oral dose of ^{14}C -BOC, the most abundant circulating metabolites were a diastereomeric mixture of ketone-reduced metabolites with a mean exposure approximately 4-fold greater than that of BOC. Boceprevir also undergoes, to a lesser extent, oxidative metabolism mediated by CYP3A4/5.

Excretion:

Boceprevir is eliminated with a mean plasma half-life ($t_{1/2}$) of approximately 3.0 hours (n = 71). Boceprevir has a mean total body clearance (CL/F) of approximately 159 L/h (n = 71). Following a single 800 mg oral dose of ^{14}C -BOC, approximately 79% and 9% of the dose was excreted in feces and urine, respectively, with approximately 8% and 3% of the dosed radiocarbon eliminated as BOC in feces and urine. The data indicate that BOC is eliminated primarily by the liver.

Ribavirin

Absorption:

Ribavirin is rapidly and extensively absorbed following oral administration. However, due to first-pass metabolism, absolute bioavailability is approximately 33% - 64%. There is a linear relationship between dose and AUC_{0-t} following single doses of 200-1,200 mg RBV. Volume of distribution is approximately 5,000 L. Based upon C_{max} from single dose to 6 weeks, accumulation of RBV in plasma is approximately 4.7 fold, although steady state may not have been achieved at 6 weeks. Following oral dosing with 600 mg RBV BID, mean plasma concentrations of 2,200 (37%) ng/mL were achieved. Upon discontinuation of dosing, the mean half-life was 298 (30%) hours, which probably reflects slow elimination from non-plasma compartments.

Ribavirin has been shown to produce high inter- and intra-subject PK variability following single oral doses (intrasubject variability of approximately 30% for both AUC and C_{max}). This may be due to extensive first pass metabolism and transfer within and beyond the blood compartment.

Distribution:

Ribavirin transport into non-plasma compartments has been most extensively studied in red cells, and has been identified to be primarily via an e_s-type equilibrative nucleoside transporter. This type of transporter is present on virtually all cell types and may account for the high volume of distribution of RBV. The ratio of whole blood:plasma RBV concentrations is approximately 60:1; the excess of RBV in whole blood exists as RBV nucleotides sequestered in erythrocytes.

Metabolism:

Ribavirin has two pathways of metabolism: (i) a reversible phosphorylation pathway in nucleated cells; and (ii) a degradative pathway involving deribosylation and amide hydrolysis to yield a triazole carboxamide metabolite. Ribavirin and its triazole carboxylic acid metabolites are excreted renally. Following oral administration of 600 mg of ¹⁴C-RBV, approximately 61% and 12% of the radioactivity was eliminated in the urine and feces, respectively, in 336 hours. Unchanged RBV accounted for 17% of the administered dose.

Effect of Food on Absorption of ribavirin: Both AUC_t (AUC from time zero to last measurable concentration) and C_{max} increased by 70% when RBV was administered with a high fat meal (841 kcal, 53.8 g fat, 31.6 g protein, and 57.4 g carbohydrate) in a single-dose PK study. It is possible that the increased bioavailability in this study was due to delayed transit of RBV or modified pH. The clinical relevance of results from this single dose study is unknown. In the pivotal clinical efficacy trial, patients were instructed to take RBV with food to achieve the maximal plasma concentration of RBV.

Peginterferon alfa-2b

Absorption:

Peginterferon alfa-2b is a well-characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species with small amounts of dipegylated and free interferon alfa-2b. The plasma half-life of PegIFNα2b is prolonged compared with non-pegylated interferon alfa-2b. Peginterferon alfa-2b has a potential to depegylate

to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar to, but lower than that of free interferon alfa-2b.

Distribution:

Following subcutaneous administration, maximal serum concentrations occur between 15 - 44 hours post-dose, and are sustained for up to 48 - 72 hours post-dose. Peginterferon alfa-2b C_{max} and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 L/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Metabolism:

Mean PegIFN α 2b elimination half-life is approximately 40 hours, with apparent clearance of 22.0 mL/hr \times kg. The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. Based on a retrospective regression analysis of PegIFN α 2b CI/F and creatinine clearance, from an expanded database, it is estimated that renal clearance of PegIFN α 2b may account for approximately 30% of the apparent clearance.

Pharmacokinetic Analysis of Combined ribavirin and peginterferon alfa-2b

Administration:

A RBV population PK analysis was conducted upon serum samples obtained at weeks 12, 24 and 48 during treatment with PEGETRON[®]. Based upon PK modeling, the recommended dose of 800/1,000/1,200 mg/day based on body weights of < 65/65 - 85/> 85 kg (in combination with PegIFN α 2b 1.5 mcg/kg), showed an overall 6.3% improved sustained response¹ rate relative to a fixed dose of 800 mg/day. The improved sustained response rate was larger (+7.4%) in the patients with HCV Genotype 1 compared to patients with HCV Genotype non-1 (3.8%).

The toxicity rate, defined as the percentage of patients with hemoglobin below 10.5 mg/dL at week four of treatment was only minimally increased by 2.5% relative to a fixed dose of 800 mg/day. This increase in toxicity was considered mild and clinically manageable.

Peginterferon alfa-2b trough concentrations were obtained at weeks 12, 24 and 48 during treatment with PEGETRON[®]. The observed concentrations and the trend toward accumulation were similar to that observed previously with PegIFN α 2b monotherapy for chronic hepatitis C, supporting the lack of PK interaction between PegIFN α 2b and RBV.

Special Populations and Conditions

Pediatrics

The safety, efficacy, and PK profile of VICTRELIS[™] in pediatric patients below the age of 18 years have not been established. Specific PK evaluations in patients under 18 years of age were not performed with PEGETRON[®]. Safety and effectiveness of PEGETRON[®] in these patients have not been evaluated. VICTRELIS TRIPLE[™] is indicated for the treatment of CHC only in patients 18 years of age or older.

¹ Sustained response was assessed by the response rate 24 weeks after the cessation of treatment

Geriatrics (≥ 65 years of age)

Population PK analysis of VICTRELIS™ indicated that age had no apparent effect on exposure.

In a single dose study using a subcutaneous dose of 1.0 mcg/kg, the pharmacokinetics of PegIFN α 2b were not affected by age. The study was not powered to detect specified differences between the age groups (20 - 45 years and 65 - 80 years). There does not appear to be a significant age-related effect on the pharmacokinetics of RBV. However, as in younger patients, renal function must be determined prior to the administration of VICTRELIS TRIPLE™.

Gender

No gender-related PK differences have been observed in adult patients that received VICTRELIS™.

Race

Population PK analysis of VICTRELIS™ indicated that race had no apparent effect on exposure.

Hepatic Insufficiency

VICTRELIS™

Boceprevir: In a study of patients with varying degrees of stable chronic liver impairment (mild, moderate and severe), no clinically significant differences in PK parameters were found and no dosage adjustment is recommended (see **DETAILED PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions, Hepatic Insufficiency**).

PEGETRON®

There is limited data available for the use of PEGETRON® in patients with mild and moderate hepatic dysfunction. In a single dose, parallel group, Phase 1 study in a limited number of patients (n = 5 to 7 patients per group) with various degrees of hepatic dysfunction (mild, moderate and severe) C_{max} increased with increasing severity of liver dysfunction ($P < 0.05$). While there were no statistically significant differences detected with AUC_t, the limited size of the study population does not permit any generalizations to be made.

Ribavirin: Single-dose pharmacokinetics of RBV in patients with mild, moderate or severe hepatic dysfunction (Child-Pugh classification A, B or C) showed a similar extent of absorption to that of normal controls.

Peginterferon alfa-2b: The pharmacokinetics of PegIFN α 2b have not been evaluated in patients with severe hepatic dysfunction. Therefore, VICTRELIS TRIPLE™ must not be used in these patients.

Renal Insufficiency

Patients with severe renal dysfunction (creatinine clearance < 50 mL/min) must not be treated with VICTRELIS TRIPLE™ (see **CONTRAINDICATIONS**).

VICTRELIS™

Boceprevir: No clinically significant differences in PK parameters were observed between patients with end-stage renal disease (ESRD) and healthy subjects (see **DETAILED PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions, Renal Insufficiency**). No dosage adjustment is required in these patients and in patients with any degree of renal impairment.

PEGETRON®

Ribavirin: The pharmacokinetics of RBV were assessed, in a limited number of subjects (n = 6 per group), after administration of a single oral dose (400 mg) of RBV to subjects with varying degrees of renal function. Both C_{max} and AUC_t of RBV appeared to increase with increasing severity of renal dysfunction. Due to the limited size of the study groups, no dosing recommendations can be made and, therefore, the use of VICTRELIS TRIPLE™ in the presence of moderate to severe renal dysfunction cannot be recommended. Single-dose RBV pharmacokinetics were altered (increased AUC_{0-24} and C_{max}) in patients with renal dysfunction compared with control subjects (creatinine clearance > 90 mL/minute). This appears to be due to reduction of apparent clearance in these patients.

Peginterferon alfa-2b: Renal clearance appears to account for 30% of total clearance of PegIFN α 2b. In a single dose study (1.0 mcg/kg) in patients with impaired renal function, C_{max} , AUC, and half-life increased in relation to the degree of renal impairment (see **CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS**). Because of marked intra-subject variability in interferon pharmacokinetics, it is recommended that patients be monitored closely during treatment with VICTRELIS TRIPLE™ (see **WARNINGS AND PRECAUTIONS**).

STORAGE AND STABILITY

Storage of VICTRELIS TRIPLE™ Packages

Store VICTRELIS TRIPLE™ package refrigerated between 2°C and 8°C. Do not use past expiry date on the label.

Storage of boceprevir

Boceprevir should be refrigerated at 2°C - 8°C until dispensed to the patient. When separated and for patient use, the product may be stored in the refrigerator until the expiration date printed on the label. The product can also be stored at room temperature (15°C - 30°C) for up to 3 months.

Store in the original container.

Storage of ribavirin

When separated, RBV should be stored in the refrigerator between 2°C and 8°C or at room temperature between 15°C and 30°C.

Storage for peginterferon alfa-2b

When separated and before reconstitution, store the individual carton of PegIFN α 2b at 2°C to 8°C. Once reconstituted PegIFN α 2b should be used immediately but may be stored at 2°C - 8°C for up to

24 hours. Do not freeze.

SPECIAL HANDLING INSTRUCTIONS

No special requirements.

Any unused product or waste material should be disposed of in accordance with local requirements.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Dosage Forms

Please refer to **DOSAGE AND ADMINISTRATION** for recommended dosing.

VICTRELIS™

Boceprevir: Boceprevir gelatin capsules are for oral administration.

PEGETRON®

Ribavirin: Ribavirin gelatin capsules are for oral administration.

Peginterferon alfa-2b: Peginterferon alfa-2b powder for solution is for subcutaneous administration.

Composition

VICTRELIS™

Boceprevir: Each hard gelatin capsule has a yellowish-brown, opaque cap with the Merck logo imprinted in red ink and off-white, opaque body with the code "314" imprinted in red ink.

Each VICTRELIS™ capsule contains 200 mg of BOC and the following inactive ingredients: croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, pre-gelatinized starch, and sodium lauryl sulfate. The capsule shell consists of gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. The capsule is printed with red ink. The red ink contains red iron oxide and shellac.

PEGETRON®

Ribavirin: Ribavirin capsules are available as opaque, white, hard gelatin capsules containing a white powder and printed with the Schering-Plough logo and "200 mg" in blue ink.

Non-medicinal ingredients are croscarmellose sodium, lactose monohydrate, magnesium stearate and microcrystalline cellulose. The capsule shell contains gelatin, sodium lauryl sulfate, silicon dioxide, and titanium dioxide.

Peginterferon alfa-2b: Peginterferon alfa-2b Powder for Solution in REDIPEN® single dose delivery system consists of a dual-chamber glass cartridge with a sterile active chamber containing PegIFN α 2b as a white to off-white lyophilized powder and a diluent chamber containing Sterile Water for Injection. The cartridge is provided in a pen device for reconstitution, dose preparation and subcutaneous administration, to deliver doses of 80, 100, 120 or 150 mcg

in 0.5 mL of reconstituted solution. Each REDIPEN[®] contains 108, 135, 162 or 202.5 mcg PegIFN α 2b. Following reconstitution of the powder with the diluent contained within the cartridge, each REDIPEN[®] gives a final volume of 0.675 mL for administration of up to 0.5 mL. The reconstituted solution contains 160, 200, 240, 300 mcg/mL respectively. Each REDIPEN[®] contains PegIFN α 2b as the active ingredient. When reconstituted as directed, each 0.5 mL solution contains 0.75 mg sodium phosphate dibasic anhydrous, 0.75 mg sodium phosphate monobasic dihydrate, 40 mg sucrose, and 0.05 mg polysorbate 80.

Packaging

VICTRELIS TRIPLE[™] is available in the following package presentations, which provide sufficient BOC, RBV and PegIFN α 2b for two weeks of VICTRELIS TRIPLE[™] therapy:

Deliverable Dose 80 mcg/0.5 mL

A carton containing two boxes of 84 BOC capsules each for a total of 168 BOC capsules, two boxes of 28 RBV capsules each for a total of 56 RBV capsules, plus two PegIFN α 2b REDIPEN[®] single dose delivery systems, 80 mcg/REDIPEN[®], with two 30-gauge needles (0.3 x 8 mm), 4 alcohol swabs and two pen holders.

Deliverable Dose 100 mcg/0.5 mL

A carton containing two boxes of 84 BOC capsules each for a total of 168 BOC capsules, two boxes of 28 RBV capsules each for a total of 56 RBV capsules, plus two PegIFN α 2b REDIPEN[®] single dose delivery systems, 100 mcg/REDIPEN[®], with two 30-gauge needles (0.3 x 8 mm), 4 alcohol swabs and two pen holders.

Deliverable Dose 120 mcg/0.5 mL

A carton containing two boxes of 84 BOC capsules each for a total of 168 BOC capsules, two boxes of 35 RBV capsules each for a total of 70 RBV capsules, plus two PegIFN α 2b REDIPEN[®] single dose delivery systems, 120 mcg/REDIPEN[®], with two 30-gauge needles (0.3 x 8 mm), 4 alcohol swabs and two pen holders.

Deliverable Dose 150 mcg/0.5 mL

1. A carton containing two boxes of 84 BOC capsules each for a total of 168 BOC capsules, two boxes of 42 RBV capsules each for a total of 84 RBV capsules, plus two PegIFN α 2b REDIPEN[®] single dose delivery systems, 150 mcg/REDIPEN[®], with two 30-gauge needles (0.3 x 8 mm), 4 alcohol swabs and two pen holders.
2. A carton containing two boxes of 84 BOC capsules each for a total of 168 BOC capsules, two boxes of 49 RBV capsules each for a total of 98 RBV capsules, plus two PegIFN α 2b REDIPEN[®] single dose delivery systems, 150 mcg/REDIPEN[®], with two 30-gauge needles (0.3 x 8 mm), 4 alcohol swabs and two pen holders.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Proper Name:

boceprevir
ribavirin
peginterferon alfa-2b

Chemical Name:

boceprevir: (1R,5S)-N-[3-Amino-1-(cyclobutylmethyl)-2,3-dioxopropyl]-3-[2(S)-[[[(1,1-dimethylethyl)amino]carbonyl]amino]-3,3-dimethyl-1-oxobutyl]-6,6-dimethyl-3-azabicyclo[3.1.0]hexan-2(S)-carboxamide

ribavirin: 1-β-D-ribofuranosyl-H-1,2,4-triazole-3-carboxamide


Molecular Formula and Molecular Mass:

boceprevir: C₂₇H₄₅N₅O₅; 519.7


ribavirin: C₈H₁₂N₄O₅; 244.2

peginterferon alfa-2b: 31,994 daltons (with a distribution from 30,000-34,000 daltons due to PEG heterogeneity)

Structural Formula:



ribavirin



peginterferon alfa-2b: Peginterferon alfa-2b is the covalent conjugate of recombinant interferon alfa-2b (IFN) with monomethoxypolyethylene glycol (PEG, average molecular weight of 12,000 daltons). Peginterferon alfa-2b is predominantly composed of monopegylated species (one PEG molecule is attached to one interferon molecule), with only a small amount of dipegylated species. Fourteen different PEG attachment sites on the interferon molecule have been identified.

Physicochemical Properties:

Boceprevir

Appearance: Boceprevir is a white to off-white amorphous powder.

Solubility: Boceprevir is freely soluble in methanol, ethanol and isopropanol and slightly soluble in water (1.5 mg/mL at 25°C).

Ribavirin

Appearance: Ribavirin is a white crystalline powder in a white opaque gelatin capsule.

Solubility: Ribavirin is freely soluble in water and slightly soluble in dehydrated alcohol.

Peginterferon alfa-2b

Appearance: Peginterferon alfa-2b Powder for Solution may appear either as a white, tablet shaped solid that is whole or in pieces, or as a white powder.

Solubility: Peginterferon alfa-2b drug substance is soluble in water and is a solution in 0.02 M sodium phosphate.

CLINICAL TRIALS

Refer to the PEGETRON[®] Product Monograph for Clinical Trial information for PEGETRON[®] specifically.

The efficacy of BOC as a treatment for CHC (genotype 1) infection was assessed in approximately 1,500 adult patients who were previously untreated (SPRINT-2) or who had failed previous therapy (RESPOND-2) in Phase 3 clinical studies.

Previously Untreated Patients (SPRINT-2)

Demographic, Disease Characteristics and Trial Design

SPRINT-2 (P05216) was conducted in previously untreated HCV, genotype 1 infected patients. A summary of the clinical trial design and patient demographics are shown in Table 13. Patients were randomized in a 1:1:1 ratio in two cohorts (Cohort 1/non-Black and Cohort 2/Black) and stratified by HCV genotype (1a or 1b) and by HCV RNA viral load ($\leq 400,000$ IU/mL vs $> 400,000$ IU/mL).

Table 13: Study P05216 (SPRINT-2) – Summary of Trial Design for Cohort 1 + Cohort 2 (all patients)

Trial Design	Dosage and Route of Administration	Treatment Regimen ^a	Total Duration (weeks)	No. of Patients	Gender M/F Race W/B/O Mean Age (years) (Range)
Phase 3, randomized, double-blinded, placebo-controlled, multi-centre	BOC – 800 mg TID, PO PegIFN α 2b – 1.5 mcg/kg/week, SC RBV – 600 to 1,400 mg/day BID (weight based dosing), PO	<u>Control</u> PegIFN α 2b/RBV–4 lead-in + (Pbo/ PegIFN α 2b/RBV–44)	Pbo – 44 PegIFN α 2b/RBV – 48	363	206/157 296/52/15 48.6 (18–75)
		<u>RGT</u> PegIFN α 2b/RBV–4 lead-in + (BOC/ PegIFN α 2b/RBV–24) ^b OR (BOC+ PegIFN α 2b/RBV–24/ PegIFN α 2b/RBV–20) ^c	BOC – 24 PegIFN α 2b/RBV – 28 OR BOC –24 PegIFN α 2b/RBV – 48	368	229/139 304/52/12 49.8 (21–76)
		<u>Not RGT</u> PegIFN α 2b/RBV–4 lead-in + (BOC/ PegIFN α 2b/RBV–44)	BOC – 44 PegIFN α 2b/RBV – 48	366	221/145 295/55/16 48.9 (21–67)

BOC = Boceprevir; PegIFN α 2b = peginterferon alfa-2b; RBV = ribavirin; Pbo = Placebo

TID = three times daily; BID = twice daily; PO = orally; SC = subcutaneous

RGT = Response-Guided Therapy (based on TW 8 results)

Race W/B/O = White/Black/Other

a: The number indicates the number of weeks of treatment

b: Early responders (Undetectable HCV-RNA at TW 8 through TW 24)

c: Late responders (Detectable HCV-RNA at TW 8 or after and undetectable at TW 24)

Futility Rule: All patients with detectable HCV-RNA in plasma at TW 24 were discontinued from treatment.

Table 14: Baseline Characteristics of Previously Untreated Patients with Chronic Hepatitis C Genotype 1 (SPRINT-2) in Cohort 1 plus Cohort 2 (all patients)

	BOC/Pegα2b/RBV (RGT) n = 368	BOC/Pegα2b/RBV-48 n = 366	Pegα2b/RBV-48 n = 363
<u>Mean Plasma HCV-RNA</u> (Log ₁₀ copies/mL)	6.52	6.53	6.54
<u>Viral Load</u> (IU/mL)			
≤ 400,000	9%	7%	7%
> 400,000	91%	93%	93%
<u>HCV Subtype</u> (Trugene) ^a			
1 (subtype unknown)	15%	13%	17%
1a	49%	51%	49%
1b	36%	36%	35%
<u>METAVIR Fibrosis</u> <u>Score</u> ^b			
F0/1/2	87%	86%	90%
F3/4	9%	11%	7%
Missing	4%	3%	3%
<u>Baseline Platelet Count</u> (10 ⁹ /L)%			
< 150	9%	10%	7%
≥ 150	91%	90%	93%
<u>Baseline ALT</u> , %			
Normal	20%	23%	26%
Elevated	80%	77%	74%
<u>Baseline Steatosis</u> ^c %			
0/1/2	94%	97%	96%
3	2%	1%	1%
Missing	4%	3%	3%

- a: HCV subtype as determined by TRUGENE HCV 5NC assay was used in subject stratification.
b: Liver histology based on central pathologist's reading. F0 = no fibrosis, F1 = portal fibrosis without septa, F2 = few septa, F3 = numerous septa without cirrhosis (bridging fibrosis), F4 = cirrhosis (advanced fibrosis)
c: Liver histology based on central pathologist's reading of the percentage cells that contain lipids. Score 0 = 0%, score 1 = > 0% and ≤ 5%, score 2 = > 5% and ≤ 32%, score 3 = > 32% and ≤ 66%.

Two percent of previously untreated patients used statins or were on opioid substitution therapy.

Study Results

The addition of BOC to PegIFN α 2b/RBV significantly increased the SVR rates compared to PegIFN α 2b/RBV alone in the combined cohort (63% to 66% BOC-containing arms vs. 38% PegIFN α 2b/RBV-48 control) for randomized patients who received at least one dose of any study medication (Full-Analysis-Set population) and decreased the length of therapy to 28 weeks for early responders (see Table 15). Overall, these SVR rates were approximately two-fold higher in patients who received the combination of BOC with PegIFN α 2b/RBV compared to the control group. Sustained Virologic Response rates for Blacks who received the combination of BOC with PegIFN α 2b/RBV were 42% to 53%; these rates are approximately two-fold higher than the SVR rate for the PegIFN α 2b/RBV-48 control (23%) (see Table 15). A secondary analysis of subjects who received at least one dose of BOC or placebo after the four-week lead-in with

PegIFN α 2b/RBV (Modified-Intent-to-Treat population) demonstrated SVR rates in the combined cohort of 67% to 68% BOC-containing arms vs. 40% PegIFN α 2b/RBV-48 control.

Table 15: Sustained Virologic Response, End of Treatment, Relapse^a and Discontinuation Rates in Previously Untreated Patients (SPRINT-2)

	FAS ^b		
	BOC/PegIFN α 2b/ RBV (RGT)	BOC/PegIFN α 2b/ RBV-48	PegIFN α 2b/RBV- 48 (Control)
Cohort 1 Plus Cohort 2 (All Patients)	n = 368	n = 366	n = 363
SVR ^c	63.3%	66.1%	37.7%
P value ^d	< 0.0001	< 0.0001	
Δ SVR	25.6	28.4	
95% CI for Δ SVR	(18.6, 32.6)	(21.4, 35.3)	
EOT^e (Undetectable HCV-RNA)	70.9%	75.7%	52.6%
Relapse	9.3%	9.1%	22.2%
Discontinuation			
During Lead-in Period	5%	3%	5%
After addition of BOC/placebo	35%	39%	54%
Discontinuation due to AEs after addition of BOC/placebo	10%	14%	12%
Treatment Failure after addition of BOC/placebo	16%	13%	34%
Completed Treatment	62%	59%	44%
Completed Follow-up	98%	98%	86%
Deaths	< 1%	< 1%	1%
Cohort 1 (Non-Black)	n = 316	n = 311	n = 311
SVR ^c	66.8%	68.5%	40.2%
P value ^d	< 0.0001	< 0.0001	
Δ SVR	26.6	28.3	
95% CI for Δ SVR	(19.1, 34.1)	(20.8, 35.8)	
EOT^e (Undetectable HCV-RNA)	74.4%	77.5%	56.6%
Relapse	9.1%	7.8%	22.8%
Discontinuation			
During Lead-in Period	4%	4%	5%
After addition of BOC/placebo	32%	36%	50%
Discontinuation due to AEs after addition of BOC/placebo	10%	14%	12%
Treatment Failure after addition of BOC/placebo	14%	11%	31%
Completed Treatment	65%	61%	48%
Completed Follow-up	97%	98%	86%
Deaths	< 1%	< 1%	1%
Cohort 2 (Black)	n = 52	n = 55	n = 52
SVR ^c	42.3%	52.7%	23.1%
P value ^d	0.0440	0.0035	
Δ SVR	19.2	29.7	
95% CI for Δ SVR	(1.6, 36.9)	(12.2, 47.1)	
EOT^e (Undetectable HCV-RNA)	50.0%	65.5%	28.8%
Relapse	12.0%	17.1%	14.3%

	FAS ^b		
	BOC/PegIFN α 2b/ RBV (RGT)	BOC/PegIFN α 2b/ RBV-48	PegIFN α 2b/RBV- 48 (Control)
Discontinuation			
During Lead-in Period	10%	0%	10%
After addition of BOC/placebo	49%	55%	77%
Discontinuation due to AEs after addition of BOC/placebo	13%	16%	15%
Treatment Failure after addition of BOC/placebo	28%	25%	53%
Completed Treatment	46%	45%	21%
Completed Follow-up	100%	98%	86%
Deaths	0%	0%	0%

- a: Relapse rate was the proportion of patients with undetectable HCV-RNA at End of Treatment (EOT) and detectable HCV-RNA at End of Follow-up (EOF) among patients who were undetectable at EOT and not missing EOF data.
- b: The Full-Analysis Set (FAS) consisted of all randomized patients (n=1,097) who received at least one dose of any study medication (PegIFN α 2b, RBV, or BOC).
- c: SVR: The last available value in the period at or after FW 24. If there is no such value, the FW 12 value is carried forward.
- d: Using the Cochran-Mantel Haenzel Chi-square test adjusted for baseline stratification factors: viral load (> 400,000 vs. \leq 400,000 IU/mL) and Genotype (1a vs. 1b).
- e: Responders at the End of Treatment.

Sustained Virologic Response Based on Lead-in Response

During the clinical studies, the lead-in phase during which PegIFN α 2b/RBV was administered for four weeks allowed for the assessment of patient interferon responsiveness immediately before the addition of BOC.

Interferon-responsiveness (as defined by $\geq 1\text{-log}_{10}$ decline in viral load at TW 4) was predictive of SVR. Boceprevir-treated patients who demonstrated interferon responsiveness by TW 4 achieved SVR rates of 81% (203/252) in BOC-RGT arm and 79% (200/254) in BOC-PR48 arm, compared to 52% (134/260) in patients treated with standard of care. Boceprevir-treated patients with $< 1\text{-log}_{10}$ decline in viral load at TW 4 (poor interferon-responsiveness) achieved SVR rates of 28% (27/97) in BOC-RGT arm and 38% (36/95) in BOC-PR48 arm compared to 4% (3/83) in patients treated with standard of care.

Sustained Virologic Response Based on TW 8 HCV-RNA Results

Response-Guided Therapy based on TW 8 response is equally effective as adding BOC to the 48-week standard of care regimen. Fifty-seven percent (208/368) of patients in the BOC-RGT arm had undetectable HCV-RNA at TW 8 (early responders). After accounting for treatment discontinuations, 44% (162/368) of patients reached TW 24 and were assigned a short (28 weeks) treatment with BOC in combination with PegIFN α 2b/RBV in the BOC-RGT arm. These BOC-RGT early responders demonstrated similar SVR rates (156/162 or 96%) after 28 weeks of treatment compared with the matched population in the BOC/PegIFN α 2b/RBV-48 arm (e.g., those patients in the BOC/PegIFN α 2b/RBV-48 arm who also had undetectable HCV-RNA at TW 8 through TW 24) (155/161 or 96%) (see Table 16).

Table 16: Sustained Virologic Response, End of Treatment and Relapse Rates in Experimental Arms with Undetectable or Detectable HCV-RNA at TW 8 through TW 24 in Previously Untreated Patients in the Combined Cohort

	FAS ^a			
	Undetectable HCV-RNA at TW 8 ^b (Early Responders)		Detectable HCV-RNA at TW 8 ^b (Late Responders)	
	BOC-RGT ^c	BOC/PegIFN α 2b/RBV-48	BOC-RGT ^c	BOC/PegIFN α 2b/RBV-48
SVR^d% (n/N)	96 (156/162)	96 (155/161)	72 (59/82)	75 (55/73)
EOT (Undetectable HCV-RNA)% (n/N)	100 (162/162)	99 (159/161)	80 (66/82)	90 (66/73)
Relapse^e% (n/N)	3 (5/161)	1 (2/157)	11 (7/66)	14 (9/64)

- a: The Full-Analysis Set (FAS) consisted of all randomized patients (n = 1,097) who received at least one dose of any study medication (PegIFN α 2b, RBV, or BOC).
- b: Per the study design, patients with undetectable HCV-RNA at TW 8 and all subsequent assays through TW 24 ended treatment at TW 28 (treatment duration assigned by Interactive Voice Response System (IVRS)).
- c: boceprevir-RGT - Patients received PegIFN α 2b/RBV for 4 weeks, then BOC 800 mg TID + PegIFN α 2b/RBV as follows: BOC 800 mg TID + PegIFN α 2b/RBV for 24 weeks (patients with undetectable HCV-RNA at TW 8 (early responders) and all subsequent assays through TW 24) or BOC 800 mg TID + PegIFN α 2b/RBV for 24 weeks followed by placebo + PegIFN α 2b/RBV for 20 weeks (patients with detectable HCV-RNA at TW 8 up to TW 24; but achieving undetectable HCV-RNA at TW 24).
- d: Sustained Virologic Response: The last available value in the period at and after FW 24. If there is no such value, the FW 12 value was carried forward.
- e: Relapse rate was the proportion of patients with undetectable HCV-RNA at EOT and detectable HCV-RNA at EOF among patients who were undetectable at EOT and not missing EOF data.

Similarly, subjects in the BOC-RGT arm with detectable HCV-RNA at any assay from TW 8 up to TW 24, but achieving undetectable HCV-RNA at TW 24 (82/368, 22%), were considered late responders and received an initial 4 weeks of PegIFN α 2b/RBV, then 24 weeks of BOC with PegIFN α 2b/RBV followed by 20 weeks of PegIFN α 2b/RBV alone in the BOC-RGT arm. These BOC-RGT late responders who were assigned to the BOC-RGT arm that received 48 weeks of treatment also had SVR rates (72%, 59/82) that were similar to those in the matched subjects in the BOC/PegIFN α 2b/RBV-48 arm (75%, 55/73) (see Table 16). These data support the concept that continued therapy with BOC in addition to PegIFN α 2b/RBV standard of care after TW 28 (as executed in the BOC/PegIFN α 2b/RBV-48 arm) does not improve SVR rates in late responders who receive a total of 48 weeks of PegIFN α 2b/RBV treatment.

Sustained Virologic Response Based on Baseline Factors

Sustained Virologic Response rates in patients in the BOC-RGT and BOC/PegIFN α 2b/RBV-48 compared to patients who received PegIFN α 2b/RBV alone with the following baseline factors were as follows: Baseline HCV-RNA > 400,000 IU/mL (62% (208/336) and 65% (220/341) vs. 34% (116/337)), advanced liver disease (F3/4) (41% (14/34) and 52% (22/42) vs. 38% (9/24)), cirrhotics (F4) (31% (5/16) and 42% (10/24) vs. 46% (6/13)), genotype 1a (59% (139/234) and 62% (147/237) vs. 34% (78/227)), and genotype 1b (71% (88/124) and 73% (85/117) vs. 40% (48/121)).

Among previously untreated patients with advanced liver disease (F3/4), the SVR rate was higher in patients who received four weeks of therapy with PegIFN α 2b/RBV followed by 44 weeks of BOC/PegIFN α 2b/RBV (52%) than in patients who received BOC-RGT (41%). However, these conclusions were based on a small sample size of patients with advanced fibrosis.

Previous Treatment Failures (RESPOND-2)

Demographic, Disease Characteristics and Trial Design

RESPOND-2 (P05101) was conducted in previously treated HCV, genotype 1 infected patients. A summary of the clinical trial design and patient demographics are shown in Tables 17 and 18. Patients were randomized in a 1:2:2 ratio and stratified based on response to their previous qualifying regimen (relapsers vs. non-responders) and by HCV subtype (1a vs. 1b).

Table 17: Study P05101 (RESPOND-2) – Summary of Trial Design

Trial Design	Dosage and Route of Administration	Treatment Regimen ^a	Total Duration (weeks)	No. of Patients	Gender M/F Race W/B/O Mean Age (years) (Range)
Phase 3, randomized, double-blinded, placebo-controlled, multi-centre	BOC – 800 mg TID, PO	<u>Control</u> PegIFN α 2b/RBV–4 lead-in + (Pbo/ PegIFN α 2b/RBV–44)	Pbo – 44 PegIFN α 2b/RBV – 48	80	58/22 67/12/1 52.9 (29–70)
		<u>RGT</u> PegIFN α 2b/RBV–4 lead-in + (BOC/ PegIFN α 2b/RBV–32) ^b	BOC – 24 PegIFN α 2b/RBV –36	162	98/64 142/18/2 52.9 (29–74)
	OR (BOC+ PegIFN α 2b/RBV–32/ PegIFN α 2b/RBV–12) ^c	OR BOC –24 PegIFN α 2b/RBV – 48			
	RBV – 600 to 1,400 mg/day BID (weight based dosing), PO	<u>Not RGT</u> PegIFN α 2b/RBV–4 lead-in + (BOC/ PegIFN α 2b/RBV–44)	BOC – 44 PegIFN α 2b/RBV – 48	161	112/49 135/19/7 52.3 (26–74)

BOC = boceprevir; PegIFN α 2b = peginterferon alfa-2b; RBV = ribavirin; Pbo = Placebo

TID = three times daily; BID = twice daily; PO = orally; SC = subcutaneous

RGT = Response-Guided therapy (based on TW 8 results)

Race W/B/O = White/Black/Other

a: The number indicates the number of weeks of treatment

b: Early responders (Undetectable HCV-RNA at TW 8)

c: Late responders (Detectable HCV-RNA at TW 8 but subsequently undetectable at TW 12)

Futility Rule: All patients with detectable HCV-RNA in plasma at TW 12 were discontinued from treatment.

Table 18: Baseline Characteristics of Previous Treatment Failures with Chronic Hepatitis C Genotype 1 (RESPOND-2)

	BOC/PegIFNα2b/RBV (RGT) n = 162	BOC/PegIFNα2b/RBV-48 n = 161	PegIFNα2b/RBV-48 n = 80
<u>Mean Plasma HCV-RNA (Log₁₀ copies/mL)</u>	6.63	6.69	6.52
<u>Viral Load (IU/mL)</u>			
≤ 400,000	4%	4%	8%
> 400,000	96%	96%	92%
<u>HCV Subtype (Trugene)^a</u>			
1 (subtype unknown)	8%	11%	8%
1a	46%	48%	48%
1b	46%	42%	45%
<u>METAVIR Fibrosis Score^b</u>			
F0/1/2	74%	72%	76%
F3/4	19%	20%	19%
Missing	7%	8%	5%
<u>Response to Qualifying Regimen</u>			
Nonresponder	35%	36%	36%
Relapser	65%	64%	64%
<u>Baseline Platelet Count (10⁹/L)%</u>			
< 150,000	13%	12%	13%
≥ 150,000	87%	88%	88%
<u>Baseline ALT,%</u>			
Normal	33%	29%	31%
Elevated	67%	71%	69%
<u>PegIFNα used in Qualifying Regimen</u>			
PEG α 2a	49%	42%	53%
PEG α 2b	51%	58%	48%
<u>Baseline Steatosis^c%</u>			
0/1/2	87%	93%	93%
3	4%	1%	1%
4	0%	0%	1%
Missing	8%	7%	5%

a: HCV subtype as determined by TRUGENE HCV 5NC assay was used in subject stratification.

b: Liver histology based on central pathologist's reading. F0=no fibrosis, F1 = portal fibrosis without septa, F2 = few septa, F3 = numerous septa without cirrhosis (bridging fibrosis), F4 = cirrhosis (advanced fibrosis)

c: Liver histology based on central pathologist's reading of the percentage cells that contain lipids. Score 0 = 0%, score 1 = > 0% and ≤ 5%, score 2 = > 5% and ≤ 32%, score 3 = > 32% and ≤ 66%, score 4 = > 66%.

Three percent of patients who failed previous therapy used statins, and one percent was on opioid substitution therapy.

Study Results

The addition of BOC to the PegIFN α 2b/RBV therapy significantly increased the SVR rates compared to PegIFN α 2b/RBV therapy alone (59% to 66% BOC-containing arms vs. 21% PegIFN α 2b/RBV-48 control) for randomized patients who received at least one dose of any study medication (Full-Analysis Set population) and decreased the length of therapy to 36 weeks for many previous treatment failures (see Table 19). Overall, these SVR rates were approximately three-fold higher in patients who received the combination of BOC with PegIFN α 2b/RBV compared to the control group. A secondary analysis of subjects who received at least one dose of BOC or placebo after the four week lead-in with PegIFN α 2b/RBV (Modified Intent to Treat population) demonstrated SVR rates of 61% to 67% in the BOC-containing arms compared to 22% PegIFN α 2b/RBV-48 control.

Table 19: Sustained Virologic Response^a, End of Treatment, and Relapse^b Rates for Previous Treatment Failures

	FAS ^c		
	BOC/PegIFN α 2b/RBV (RGT) n = 162	BOC/PegIFN α 2b/RBV-48 n = 161	PegIFN α 2b/RBV-48 (Control) n = 80
SVR ^a	58.6%	66.5%	21.3%
P value ^d	< 0.0001	< 0.0001	
Δ SVR	37.4	45.2	
95% CI for Δ SVR	(25.7, 49.1)	(33.7, 56.8)	
EOT (Undetectable HCV-RNA)	70.4%	77.0%	31.3%
Relapse	15.3%	11.6%	32.0%
Discontinuation			
During Lead-in Period	4%	1%	3%
After addition of BOC/placebo	33%	34%	71%
Discontinuation due to AEs after addition of BOC/placebo	6%	12%	1%
Treatment Failure after addition of BOC/placebo	23%	18%	63%
Completed Treatment	64%	65%	29%
Completed Follow-up	97%	96%	97%
Deaths	< 1%	0	0

- a: Sustained Virologic Response: The last available value in the period at and after FW 24. If there is no such value, the FW 12 value was carried forward.
- b: Relapse rate was the proportion of patients with undetectable HCV-RNA at EOT and detectable HCV-RNA at EOF among patients who were undetectable at EOT and not missing EOF data.
- c: The FAS consisted of all randomized patients (n = 403) who received at least one dose of any study medication (PegIFN α 2b, RBV, or BOC).
- d: Using the Cochran-Mantel Haenzel Chi-square test adjusted for the baseline stratification factors: prior response status (some negative vs. never negative) and Genotype (1a vs. 1b).

Sustained Virologic Response Based on Previous Treatment Response and Lead-in Response
Achievement of SVR was associated with the patient's response to PegIFN α 2b/RBV therapy, whether defined by classification of response to previous treatment, or by a decrease in HCV-RNA at TW 4 (see Table 20). The TW 4 response was a stronger predictor of SVR compared to response to previous treatment and allowed the determination of the patient's on-treatment interferon responsiveness.

Table 20: Sustained Virologic Response, End of Treatment, Relapse^a Rates for Previous Treatment Failures by Previous Treatment Response and Lead-In Response (FAS^b)

	Previous Treatment Response		Lead-In Response ^c (TW 4) HCV-RNA Log Reduction	
	Previous Non-Responders ^c	Previous Relapsers ^d	< 1- \log_{10} decline	\geq 1- \log_{10} decline
PegIFNα2b/RBV-48 (n = 80)				
SVR ^f % (n/N)	7 (2/29)	29 (15/51)	0 (0/12)	25 (17/67)
Relapse ^a % (n/N)	33 (1/3)	32 (7/22)	0 (0/0)	32 (8/25)
EOT% (n/N)	10 (3/29)	43 (22/51)	0 (0/12)	37 (25/67)
BOC-RGT (n = 162)				
SVR ^f % (n/N)	40 (23/57)	69 (72/105)	33 (15/46)	73 (80/110)
Relapse ^a % (n/N)	18 (5/28)	14 (12/83)	12 (2/17)	16 (15/94)
EOT% (n/N)	54 (31/57)	79 (83/105)	41 (19/46)	86 (95/110)
BOC/PegIFNα2b/RBV-48 (n = 161)				
SVR ^f % (n/N)	52 (30/58)	75 (77/103)	34 (15/44)	79 (90/114)
Relapse ^a % (n/N)	14 (5/35)	10 (9/86)	25 (5/20)	9 (9/99)
EOT% (n/N)	60 (35/58)	86 (89/103)	48 (21/44)	89 (101/114)

- a: Relapse rate was the proportion of patients with undetectable HCV-RNA at EOT and detectable HCV-RNA at EOF among patients who were undetectable at EOT and not missing EOF data.
- b: The FAS consisted of all randomized patients (n = 403) who received at least one dose of any study medication (PegIFN α 2b, RBV, or BOC).
- c: Previous Non-Responder = patient who failed to achieve SVR after at least 12 weeks of previous treatment with PegIFN α 2b/RBV, but demonstrated a \geq 2- \log_{10} reduction in HCV-RNA by Week 12.
- d: Previous Relapser = patient who failed to achieve SVR after at least 12 weeks of previous treatment with PegIFN α 2b/RBV, but had undetectable HCV-RNA at the EOT.
- e: Eleven patients were missing TW 4 assessment (HCV-RNA) and were not included in the Lead-In response results.
- f: Sustained Virologic Response: The last available value in the period at and after FW 24. If there is no such value, the FW 12 value was carried forward.

Sustained Virologic Response Based on TW 8 HCV-RNA Results

Response-Guided Therapy based on TW 8 response is equally effective as adding BOC to the 48-week standard of care regimen. Forty-six percent (74/162) of patients in the BOC-RGT arm and 52% (84/161) of patients in the BOC/PegIFN α 2b/RBV-48 arm were early responders (patients with undetectable HCV-RNA at TW 8). Of the patients that were early responders, 71 patients were undetectable at TW 12 in the BOC-RGT arm and 81 patients were undetectable at TW 12 in BOC/PegIFN α 2b/RBV-48 arm. Boceprevir-RGT early responders, who received 36 weeks of therapy (an initial 4 weeks of PegIFN α 2b/RBV followed by 32 weeks of BOC with PegIFN α 2b/RBV), had an SVR rate of 86% (64/74) compared with an SVR rate of 88% (74/84) in the matched population in the BOC/PegIFN α 2b/RBV-48 arm who received 48 weeks of therapy (an initial 4 weeks of PegIFN α 2b/RBV followed by 44 weeks of BOC with PegIFN α 2b/RBV) (see Table 21).

Table 21: Sustained Virologic Response, End of Treatment, and Relapse Rates in the Experimental Arms with Undetectable or Detectable HCV-RNA at TW 8 in Previous Treatment Failures

	Undetectable HCV-RNA at TW 8		Detectable HCV-RNA at TW 8	
	BOC-RGT ^a	BOC/PegIFN α 2b/RBV-48	BOC -RGT ^{a,b}	BOC/PegIFN α 2b/RBV-48
SVR ^c %, (n/N)	86 (64/74)	88 (74/84)	40 (29/72)	43 (30/70)
EOT (Undetectable HCV-RNA)% (n/N)	97 (72/74)	96 (81/84)	56 (40/72)	57 (40/70)
Relapse ^d % (n/N)	11 (8/71)	8 (6/80)	24 (9/38)	21 (8/38)

- a: BOC-RGT - Patients received PegIFN α 2b/RBV for 4 weeks, then BOC 800 mg TID + PegIFN α 2b/RBV as follows: BOC 800 mg TID + PegIFN α 2b/RBV for 32 weeks (patients with undetectable HCV-RNA at TW 8 (early responders) and TW 12) or BOC 800 mg TID + PegIFN α 2b/RBV for 32 weeks followed by placebo + PegIFN α 2b/RBV for 12 weeks (patients detectable HCV-RNA at TW 8 but subsequently negative by TW 12).
- b: Sustained Virologic Response: The last available value in the period at and after FW 24. If there is no such value, the FW 12 value was carried forward.
- c: Relapse rate was the proportion of patients with undetectable HCV-RNA at EOT and detectable HCV-RNA at EOF among patients who were undetectable at EOT and not missing EOF data.

In patients who were not early responders (patients with detectable HCV-RNA at TW 8), the SVR rate in the BOC-RGT arm was 40% (29/72) compared with an SVR rate of 43% (30/70) in the matched population in the BOC/PegIFN α 2b/RBV-48 arm (see Table 21). Thirty-eight patients in the BOC-RGT arm and 37 patients in the BOC/PegIFN α 2b/RBV-48 arm had detectable HCV-RNA at TW 8 but were subsequently undetectable at TW 12 (late responders). Boceprevir-RGT late responders, who received an initial 4 weeks of PegIFN α 2b/RBV then 32 weeks of BOC with PegIFN α 2b/RBV followed by 12 weeks of PegIFN α 2b/RBV alone, had an SVR rate of 76% (29/38) compared with an SVR rate of 62% (23/37) in the matched population in the BOC/PegIFN α 2b/RBV-48 arm, who received 4 weeks of PegIFN α 2b/RBV followed by 44 weeks of BOC in addition to PegIFN α 2b/RBV. These data support that, in late responders, 36 weeks of BOC with PegIFN α 2b/RBV followed by 12 weeks of PegIFN α 2b/RBV is adequate and that treatment with BOC may be shortened to 32 weeks in patients who have received previous therapy.

A difference was observed in the number of patients who achieved SVR between the BOC-RGT arm and the BOC/PegIFN α 2b/RBV-48 arm. This difference is explained by imbalances in treatment response observed amongst patients in each arm who received identical therapy prior to TW 36.

Sustained Virologic Response Based on Baseline Factors

Sustained Virologic Response rates of patients in the BOC-RGT and BOC/PegIFN α 2b/RBV-48 compared to patients who received PegIFN α 2b/RBV alone with the following baseline factors were as follows: Baseline HCV-RNA > 400,000 IU/mL (57% (88/155) and 66% (102/154) vs. 19% (14/74)), advanced liver disease (F3/4) (44% (14/32) and 68% (21/31) vs. 13% (2/15)), cirrhotics (F4) (35% (6/17) and 77% (17/22) vs. 0% (0/10)), genotype 1a (53% (50/94) and 64% (61/96) vs. 24% (11/46)), and genotype 1b (67% (44/66) and 70% (43/61) vs. 18% (6/34)).

Among patients who failed previous therapy with advanced liver disease (F3/4), the SVR rate was higher in patients who received four weeks of therapy with PegIFN α 2b/RBV followed by 44 weeks of BOC/PegIFN α 2b/RBV (68%) compared to patients who received BOC-RGT (44%). However, these conclusions were based on a small sample size of patients with advanced fibrosis.

DETAILED PHARMACOLOGY

Pharmacodynamics

Boceprevir

Electrocardiogram (ECG) Evaluation

Thirty-six healthy adult subjects were enrolled into and 31 subjects completed a randomized, placebo- and active-controlled, multiple-dose, evaluator-blind, four-way, crossover study. Each treatment consisted of oral dosing for 5 days of: A) BOC 800 mg TID; B) BOC 1,200 mg TID; C) moxifloxacin 400 mg Once Daily (QD); and D) placebo with a 7-day washout period between each treatment period. The moxifloxacin treatment group validated the study design since prolongation of mean QT/QTc was measured with this treatment. Boceprevir did not prolong the QTcF interval at the 800 mg (therapeutic dose) or the 1,200 mg dose or relative to placebo. Neither the 800 mg dose or the 1,200 mg dose of BOC was associated with clinically relevant effects on cardiac conduction.

At the mean maximum BOC exposures of 1,690 and 1,940 ng/mL, which were achieved at 2 hours postdose for the 800 and 1,200 mg doses, the mean increases in placebo-adjusted QTcF were 4.5 and 0.3 ms with the upper limits of the 95% CI of 7.3 and 3.1 ms, respectively. The maximum observed mean increases in placebo-adjusted QTcF occurred at 4 hours post-dose, and were 5.8 and 2.9 ms with the upper limits of the 95% CI of 8.7 and 5.7 ms, respectively.

Therefore, in this study with demonstrated ability to detect small effects, the upper bound of the one-sided 95% confidence interval for the largest placebo-adjusted, baseline-corrected QTc based on individual correction method was below 10 ms, the threshold for regulatory concern. The dose of 1,200 mg yields a BOC maximum exposure increase of approximately 15% which may not cover exposures due to co-administration with strong CYP3A4 inhibitors or use in patients with severe hepatic impairment. However, at the doses studied in the thorough QT study, no apparent

concentration-QT relationship was identified. Thus, there is no expectation of a QTc effect under a higher exposure scenario.

When QTcF in men and women was analyzed separately, similar results were obtained. There was no difference between men and women when placebo and BOC at both the 800 mg dose and the 1,200 mg dose were compared.

Ribavirin

Ribavirin is a synthetic nucleoside analog, which has shown *in vitro* activity against some, but not all, RNA and DNA viruses. Neither RBV nor its intracellular nucleotide metabolites at physiologic concentrations have been shown to inhibit HCV-specific enzymes or HCV replication. Ribavirin is not incorporated into either RNA or DNA, and RBV treatment per se does not induce endogenous synthesis of alpha interferons.

Peginterferon alfa-2b

In a rising single-dose study, interferon pharmacodynamics were assessed by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), and white cell and neutrophil counts. Patients treated with PegIFN α 2b showed mild dose-related elevations in body temperature with the maximum temperature and change (38.0°C and 2.0°C, respectively) occurring in the 0.7 mcg/kg dose group. In comparison, patients treated with interferon alfa-2b (10 MIU) had average maximum body temperature and change from baseline values of 38.6°C and 2.7°C, respectively.

Peginterferon alfa-2b increased serum neopterin concentrations in a dose-dependent manner, and also increased 2'5'-OAS concentrations (although with no clear dose-relationship). Interferon alfa-2b increased concentrations of both effector proteins. Similar effects of PegIFN α 2b and interferon alfa-2b on hematological variables (white cell, neutrophil and platelet counts) were likewise observed. In the rising multiple-dose study, reductions in serum HCV-RNA levels were observed after administration of PegIFN α 2b.

Ribavirin plus peginterferon alfa-2b

Studies to date have not elucidated a mechanism of antiviral synergy between RBV and PegIFN α 2b that explains the increased efficacy of the combination in the treatment of chronic hepatitis C relative to either treatment alone.

Pharmacokinetics

Boceprevir

General Pharmacokinetic Characteristics

The PK profile of BOC in healthy subjects, HCV-infected patients and population PK parameters are presented in Table 22. In general, the PK results were similar between healthy subjects and HCV patients.

Table 22: Pharmacokinetics of boceprevir in Healthy Subjects, HCV-Infected Patients and PPK Estimates

Pharmacokinetic Parameters	HCV Patients (800 mg) ^a	Population PK Estimates (HCV Patients; 800 mg)	Healthy Subjects (800 mg) ^b
C _{max} (ng/mL)	1,013	1,084	1,723
AUC (ng•h/mL)	4,403	4,642	5,408
C _{min} (ng/mL)	213	218	88
T _{1/2} (h)	---	---	3
T _{max} (h)	2	1.90	2
CL/F (L/h)	182	172	159
Vc/F	207	196	---
Vd/F (L)	---	---	717

a: PPK individual prediction from sparse data

b: Parameters obtained using non-compartmental analysis

Absorption

Effects of Food on Oral Absorption

The PK parameters of BOC following administration under fasted and fed (high-fat) conditions in healthy subjects are presented in Table 23. Following administration with food to healthy subjects, BOC was rapidly absorbed with a median T_{max} of approximately 2.0 hours. *In vitro*, BOC has been shown to be a substrate of P-gp. There was no discernible dose-related effect on T_{max}. Steady-state mean BOC AUC, C_{max}, and C_{min} increased in a less than dose proportional manner and individual exposures overlapped substantially at 800 mg and 1,200 mg, suggesting diminished absorption at higher doses. The power log model shows an increase in exposure of 27% to 38% between 800 mg and 1,200 mg TID. Steady state was reached after approximately 1 day of TID dosing.

Table 23: Pharmacokinetic Parameters of boceprevir following Administration under Fasted and Fed (High-Fat) Conditions in Healthy Adult Subjects

Pharmacokinetic Parameters	Fasted	Fed (High-Fat Meal)	Ratio Estimate (%)	Confidence Interval (90%)
AUC (ng•hr/mL)	4,210	6,350	161	131–198
C _{max} (ng/mL)	1,370	1,710	148	102–216
T _{max} (hr)	1.5	3.5	---	---
t _{1/2} (hr)	3.58	3	---	---

Administration with food increased the oral bioavailability of BOC relative to the fasted state, by 40% to 60% based on AUC. Administration with food also generally modestly to delayed median BOC T_{max} from approximately 1 hour to approximately 2 hours. Meal type, and timing of meal administration, did not notably affect the increase in exposure.

Distribution

Mean Vd/F in fed subjects showed extensive distribution with a mean Vd/F of approximately 717 L (n = 71). Boceprevir is not highly bound to human plasma proteins. The mean unbound fraction of BOC in plasma was similar between healthy subjects and subjects with End-Stage Renal Disease (ESRD).

Metabolism

Clinical data show that BOC is rapidly metabolized to an inactive, ketone-reduced metabolite. This metabolite has been shown to be inactive as a serine PI in an *in vitro* assay using recombinant HCV NS3/4A protease. The median T_{\max} for this metabolite was approximately 3 hours. This metabolite is present in plasma in a ratio to BOC of approximately 4:1 at a dose of 800 mg. Based on all known *in vitro* data and clinical drug-drug interactions studies; metabolism clinically appears to be primarily via the AKR enzymes and to a minor extent via CYP3A4/5-mediated oxidation. Profiling of drug-derived material in plasma, urine and feces showed that ^{14}C -BOC was extensively metabolized by humans. Over sixty metabolites were detected in humans. The metabolic modification can be assigned to one or more of the five regions in the molecule. The major biotransformation pathway involved reduction of the second carbonyl group at the carboxamide terminus, which accounted for at least ~22% of the dose excreted in urine and feces. Other metabolic pathways included oxidation, cleavage, dimerization, and a combination of these processes, including reduction. Most of the oxidative metabolites were excreted in feces. Approximately 8% of the dose was detected in feces as unchanged drug.

Excretion

Boceprevir was eliminated with a mean plasma $t_{1/2}$ of approximately 3.0 hours ($n = 71$). The mean $t_{1/2}$ tended to be variable between studies and was associated with a high coefficient of variation (% CV) of 90%. The mean CL/F of BOC across several studies was approximately 159 L/h. Accumulation of BOC was minimal with multiple days of TID dosing and steady state was reached after approximately 1 day of TID dosing. In a clinical study with ^{14}C -BOC a mean total of 88.2% of the radioactive dose was recovered in the urine and feces 168 hours after a single oral administration of 800 mg. Radioactivity recovered in the urine and feces accounted for 9.28% and 78.9% of the dose, respectively with approximately 3% and 8% of the dosed radiocarbon eliminated as BOC in urine and feces, respectively. Most of the radioactivity was excreted in the urine within the first 12 hours.

Drug-Drug Interactions

Boceprevir is a strong inhibitor of CYP3A4/5. Drugs metabolized primarily by CYP3A4/5 may have increased exposure when administered with BOC, which could increase or prolong their therapeutic and adverse effects (see **CONTRAINDICATIONS**). Boceprevir does not inhibit CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 or CYP2E1 *in vitro*. In addition, BOC does not induce CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19 or CYP3A4/5 *in vitro*.

Boceprevir is primarily metabolized by AKR. In drug interaction trials conducted with AKR inhibitors diflunisal and ibuprofen, BOC exposure did not increase to a clinically significant extent. Boceprevir may be co-administered with AKR inhibitors.

Boceprevir is partly metabolized by CYP3A4/5. It is also a substrate for P-glycoprotein (P-gp). Co-administration of BOC with drugs that induce or inhibit CYP3A4/5 could decrease or increase exposure to BOC. Drugs that are potent CYP3A4/5 inducers may significantly reduce BOC plasma concentrations, which may be associated with reduced efficacy (see **CONTRAINDICATIONS**).

Drug interaction studies were performed with BOC and drugs most likely to be co-administered with BOC. The effects of co-administration of other drugs on the PK (AUC, C_{max} and C_{min}) of BOC are summarized in Table 24 while the effects of BOC on other drugs are summarized in Table 25. For information regarding clinical recommendations, see **DRUG INTERACTIONS**.

Table 24: Drug Interactions; PK Parameters for boceprevir in the Presence of Co-administered Drugs

	Dose/Schedule		n	Change in PK	Ratio Estimate (90% CI) of BOC PK Parameters With/Without Co-administered Drug. No Effect=1.00		
	Co-administered Drug	BOC			C _{max}	AUC	C _{min}
Co-Administration with Antifungals							
ketoconazole	400 mg BID	400 mg Single dose	12	↑	1.41 (1.00–1.97)	2.31 (2.00–2.67)	N/A
Co-Administration with Other Antivirals							
efavirenz	600 mg daily	800 mg TID	12	↑	0.92 (0.78–1.08)	0.81 (0.75–0.89)	0.56 (0.42–0.74)
ritonavir	100 mg daily	400 mg TID	12	↓	0.73 (0.57–0.93)	0.81 (0.73–0.91)	1.04 (0.62–1.75)
tenofovir	300 mg daily	800 mg TID	17	↔	1.05 (0.98–1.12)	1.08 (1.02–1.14)	1.08 (0.97–1.20)
PegIFN α 2b	1.5 mcg/kg SC weekly	400 mg TID	10	↔	0.88 (0.66–1.18)	1.00 (0.89–1.13)	N/A
Co-Administration with Other Drugs							
ibuprofen	600 mg TID	400 mg Single dose	12	↔	0.94 (0.67–1.32)	1.04 (0.90–1.20)	N/A
diflunisal	250 mg BID	800 mg TID	12	↔	0.86 (0.56–1.32)	0.96 (0.79–1.17)	1.31 (1.04–1.65)

SC = subcutaneous

Table 25: Drug Interactions; PK Parameters of Co-administered Drugs in the Presence of boceprevir

	Dose/Schedule		n	Change in PK	Ratio Estimate (90 or 95% CI) of Co-administered Drug PK Parameters With/Without BOC. No Effect = 1.00	
	Co-administered Drug	BOC			C _{max}	AUC
Co-Administration with Sedative/Hypnotics						
midazolam	4 mg single oral dose	800 mg TID	12	↑	5.30 (4.66–6.03)	2.77 (2.36–3.25)
Co-Administration with Oral Contraceptives						
drospirenone/ethinyl estradiol	drospirenone: 3 mg daily ethinyl estradiol: 0.02 mg daily	800 mg TID	16	↑ ↓	drospirenone: 1.99 (1.87–2.11) ethinyl estradiol: 0.76 (0.73–0.79)	drospirenone: 1.57 (1.46–1.70) ethinyl estradiol: 1.00 (0.91–1.10)
Co-Administration with Other Antivirals						
tenofovir	300 mg daily	800 mg TID	17	↔	1.05 (1.01–1.09)	1.32 (1.19–1.45)
efavirenz	600 mg daily	800 mg TID	12	↑	1.20 (1.15–1.26)	1.11 (1.02–1.20)
PegIFNα2b	1.5 mcg/kg SC weekly	400 mg TID	10	↔	0.99 ^{a, b} (0.83–1.17)	N/A

a: 0–168 hours

b: Reported AUC is 200 mg and 400 mg cohorts combined.

SC = subcutaneous

Population Pharmacokinetics

Population PK analysis in Phase 3 studies systemically checked for significant effects on systemic clearance, volume of distribution and absorption rates by the following characteristics: health status (HCV subject vs healthy subject), demographics (sex, black race, Asian race, age, weight, and BMI), hepatic function (AST, ALT), and renal function (creatinine clearance). Systemic clearance and volume of distribution were not influenced by demographics, measures of renal function or hepatic function, except for sex on clearance and absorption rate and of health status on central volume. These effects were well within the range of estimated inter-individual and intra-individual variability in BOC exposure, and were therefore not considered clinically relevant.

Special Populations and Conditions

Sex, Age, Body Weight, Height, and Body Mass Index

In the Phase 2 PPK analysis, sex, body weight, height, and BMI had no significant effect on BOC CL/F or Vd/F; however, an age effect was observed against CL/F. The estimated age effect on CL/F was -0.291 suggesting minimal difference in CL/F attributable to age. In the Phase 3 PPK analysis, no age effect was noted. However, effects of sex on clearance and absorption rate were noted. These effects were well within the range of estimated inter-individual and intra-individual variability in BOC exposure, as well as within clinical bounds of comparability, and were therefore not considered clinically relevant.

Race

At the target clinical dose of 800 mg TID (with food), BOC was administered for seven days to six paired healthy Caucasian and Japanese subjects (matched for age, weight and height). In general, the PK parameters of BOC following single and multiple doses of BOC were similar between Caucasian and Japanese subjects, under both fed and fasting conditions, with no notable accumulation of BOC in plasma and similar CL/F between Caucasian and Japanese subjects. At the target clinical dose of 800 mg TID administered with food, steady-state mean AUC and C_{max} in Japanese were 14% and 5% lower, respectively compared with matched-Caucasian subjects and are well within the apparent lower bounds of clinical significance. Dose-normalized pooled data analysis indicated that both the steady-state mean AUC and C_{max} of BOC increased in a less than dose-proportional manner in both Japanese and Caucasian subjects after receiving multiple doses of 200 mg, 400 mg, and 800 mg. Food increased exposure of BOC compared with fasted conditions in both Caucasian and Japanese subjects, with the increased exposure more apparent at higher dose levels. The mean ratio estimate for AUC at a single dose of 800 mg ranged from 142% to 196% fed vs. fasted. Boceprevir should be administered with food, and no modification of BOC dose is required for persons of Japanese descent. In a Phase 1 study, mean AUC and C_{max} were lower in Black compared with non-Black subjects. However, the sample size in this study was very small. The Phase 2 PPK analysis revealed that race had no significant effect on BOC CL/F or Vd/F. The non-Caucasian subjects in this subanalysis were not specific for Black subjects as there were too few Blacks, but covered all non-Caucasian races. The Phase 3 PPK analysis indicated that Black or Asian race was not a significant covariate for BOC PK.

Hepatic Insufficiency

A study was performed to assess the safety and PK of BOC in subjects with varying degrees of hepatic insufficiency. Adult non HCV-infected males and females with mild (Child-Pugh score 5-6), moderate (Child-Pugh score 7-9), severe (Child-Pugh score 10-12) impairment and matched subjects with normal hepatic function were administered a single 400 mg dose of BOC in the original formulation under fasted conditions. With increasing severity of liver impairment, a trend toward increased mean area under the plasma concentration-time curve from last quantifiable sample ($AUC_{(tf)}$) and mean C_{max} of BOC was observed (see Table 26). Mean CL/F values in subjects with moderate and severe hepatic impairment were decreased. However, mean CL/F values remained in the range of mean CL/F seen in healthy subjects from other studies. Certain conditions of this trial, fasted dosing, a less than therapeutic dose, non-final formulation, limited the generalizability of the conclusions.

Table 26: PK Parameters of AUC_(tf) and C_{max} of Hepatically Impaired Subjects Compared to Healthy Subjects for the boceprevir Active Diastereoisomer 534128

PK Parameter	Group	n	LS Mean	Treatment Comparison	Ratio Estimate%	90% CI
AUC _(tf) (ng*hr/mL)	Mild	6	1,009	Mild vs. Healthy	107	75–152
	Moderate	6	1,240	Moderate vs. Healthy	132	93–187
	Severe	6	1,361	Severe vs. Healthy	145	102–205
	Healthy	6	941	--	--	--
C _{max} (ng/mL)	Mild	6	295	Mild vs. Healthy	115	71–188
	Moderate	6	327	Moderate vs. Healthy	128	79–208
	Severe	6	413	Severe vs. Healthy	162	99–263
	Healthy	6	256	--	--	--

Estimates of steady-state maximum AUC and C_{max} parameters of patients infected with HCV in the Phase 3 studies were 9,715 ng·h/mL and 2,377 ng/mL, respectively. The Phase 3 PPK analysis showed that hepatic dysfunction was not a significant covariate for BOC PK. Based on these data, no dose adjustment of BOC is required for subjects with hepatic impairment.

However, because PegIFNα2b/RBV is contraindicated in the hepatically impaired population, the use of BOC with PegIFNα2b/RBV is also contraindicated in this population.

Renal Insufficiency

In a study with ¹⁴C-BOC, drug-derived radiocarbon was mainly eliminated via the feces, with < 10% recovered in urine, indicating renal clearance is a minor pathway. A study was performed to assess the safety, tolerability and PK of BOC in subjects with varying degrees of renal insufficiency. Male and female, non HCV-infected hemodialysis-dependent ESRD subjects and matched subjects with normal renal function were administered a single 800 mg dose of BOC. ESRD subjects were dosed prior to dialysis (Day 1) and 4 hours prior to dialysis (Day 4). The difference in exposure compared with healthy subjects was not clinically relevant, and dialysis did not affect the PK of BOC (see Table 27). As there were no differences in exposure in the most severely renally impaired subjects, no dose adjustment of BOC is required in patients with any degree of renal impairment.

Table 27: Mean (CV, %) Plasma PK Parameters of boceprevir following a Single Oral Dose of boceprevir 800 mg to Healthy Subjects and to Patients with ESRD

Parameter	Healthy Subjects	ESRD Patients	
		Day 1	Day 4
AUC _(tf) (ng·hr/mL)	5,710 (50)	5,100 (53)	5,000 (43)
AUC _(I) (ng·hr/mL)	5,760 (50)	5,150 (53)	5,030 (43)
C _{max} (ng/mL)	1,730 (54)	1,340 (52)	1,420 (35)
T _{max} ^a (hr)	2.00 (2.00-4.00)	4.00 (1.00-6.00)	2.00 (1.32-2.00)
t _{1/2} (hr)	1.73 (21)	2.20 (60)	1.72 (43)
CL/F (L/hr)	178 (55)	193 (50)	183 (38)
Vd/F (L)	417 (47)	637 (89)	495 (86)

a: Median (range)

Animal Pharmacology

Single-dose administration of ^{14}C -BOC under fasted conditions showed that concentrations of radioactivity in plasma were greater than those in whole blood, and the comparison of radioactivity concentrations in plasma and blood indicated a minor partitioning into the cellular components of blood. Mean exposure to BOC in plasma accounted for approximately 26.2% of the total plasma radioactivity. Individual plasma to blood radioactivity concentration ratios remained constant over the quantifiable intervals following dosing. Tissue distribution was examined by quantitative-whole-body autoradiography in rats administered a single, oral dose of 25 mg/kg ^{14}C -BOC.

Peak radiocarbon concentrations were observed in blood and most tissues at 0.5 hours postdose and declined to below quantifiable limits by 24 hours postdose. The highest radiocarbon concentrations were measured in liver, bladder wall, kidneys, and prostate gland. ^{14}C -BOC -derived radiocarbon was not detected in the brain or spinal cord. There were no qualitative differences in tissue distribution between male and female or between pigmented and non-pigmented rats. In preclinical studies, the percent binding increased as the concentration of BOC decreased in plasma from several species. In studies in rats, dogs, and monkeys administered ^{14}C -BOC, the main route of elimination was also via the feces, due to a combination of biliary excretion and, for orally dosed animals, unabsorbed drug. Less than 10% of administered radiocarbon was recovered in urine. Absolute bioavailability ranged from 26% to 34% in fasted mice, rats, and dogs and was 4% in monkeys. On average 71% of the excreted radioactivity in feces was due to metabolites.

Safety pharmacology studies were conducted to evaluate the effect of BOC on the cardiovascular, respiratory, central nervous, gastrointestinal and renal systems. No effects on cardiac function or electrocardiography parameters were observed in dogs with oral doses of 3 or 50 mg/kg BOC. The corrected QT interval (QTc) revealed no significant difference between BOC (75 or 200 mg/kg) and vehicle in cynomolgus monkeys. There were no statistically significant changes in respiratory rate, tidal volume or minute volume with single oral gavage doses of 25, 75 or 200 mg/kg of BOC in rats. Boceprevir did not demonstrate any drug-related effect on renal function, gastric emptying or intestinal transit. The NOAEL for central nervous system pharmacological activity in rats is 200 mg/kg based on the lack of BOC-related changes.

Ribavirin

Ribavirin is rapidly absorbed following oral administration (mean T_{\max} = 1.5 hours), followed by rapid distribution and prolonged elimination phases (single dose half-lives of absorption, distribution and elimination are 0.05, 3.73 and 79 hours respectively). Absorption is extensive, with approximately 10% of a radiolabeled dose excreted in the feces. However, absolute bioavailability is approximately 33-64%, due to first pass metabolism. Apparent volume of distribution is approximately 5,000 L, which reflects the extensive distribution of RBV.

Ribavirin accumulation in non-plasma compartments has been most extensively studied in red cells, and has been identified as primarily via an e_s -type equilibrative nucleoside transporter. This type of transporter is present on virtually all cell types, and may account for the high volume of distribution. Ribavirin has two pathways of metabolism: (i) a reversible phosphorylation pathway; and (ii) a degradative pathway involving deribosylation and amide hydrolysis to yield a

triazole carboxamide metabolite. Both RBV and its triazole carboxamide and triazole carboxylic acid metabolites are also excreted renally. Ribavirin does not bind to plasma proteins.

Based upon C_{max} from single dose to 6 weeks, accumulation of RBV in plasma is approximately 4.7 fold, although steady state may not have been achieved at 6 weeks. Following oral dosing with 600 mg RBV BID, mean plasma concentrations of 2,200 (37%) ng/mL were achieved. Upon discontinuation of dosing there was a washout half-life of approximately 298 hours, which probably reflects slow elimination from non-plasma compartments.

Population PK analysis was performed using sparsely sampled serum concentration values from the clinical efficacy studies. The clearance model developed showed that body weight, gender, age and serum creatinine were the main covariates. The estimates of RBV clearance developed by the final model were 21.1 L/hr for males, and 17.7 L/hr for females. These estimates are close to those obtained from intensively sampled multiple dose plasma PK data (approximately 23 L/hr). For males, clearance was approximately 20% higher than for females. Clearance increased as a function of body weight and was reduced at ages greater than 40 years. Effects of covariates on RBV clearance appear to be of limited clinical significance, due to the substantial residual variability not accounted for by the model. Comparison of concentration data with pharmacodynamic variables showed a positive relationship with nadir hemoglobin values or percent change in hemoglobin from baseline, although these data were highly variable. Mean concentration values between treatment responders and non-responders were generally similar, although these data were also highly variable.

The bioavailability of RBV was increased by coadministration of a high fat meal (AUC_t and C_{max} both increased by 70%).

Ribavirin is not a substrate for, nor does it inhibit or induce any cytochrome P450 enzymes. The pharmacokinetics of RBV was not assessed in elderly or pediatric subjects.

Peginterferon alfa-2b

The PK profile of PegIFN α 2b following subcutaneous injection, based on a single study, is summarized in Table 28 below. In addition, the pharmacokinetics of non-pegylated interferon alfa-2b is described for comparative purposes.

Table 28: Mean (% CV) Pharmacokinetic Parameters in Humans following Single Dose Administration of peginterferon alfa-2b

Compound	Dose (mcg/kg)	C _{max} (pg/mL)	T _{max} (hr)	AUC _(0-t) (pg×hr/mL)	t _{1/2} (hr)	Cl/F (mL/hr×kg)	Vd/F (L/kg)
PegIFNα2b^a	0.5	295 (26)	37 (46)	16,000 (32)	27.2 ^b	23.0 ^b	0.89 ^b
	1	554 (38)	31 (45)	41,400 (25)	33.4 (33)	21.9 (17)	1.07 (39)
	1.5	785 (47)	44 (22)	63,300 (43)	28.2 (27)	25.0 (40)	1.08 (57)
	2	1,710 (39)	15 (35)	105,000 (26)	31.6 (17)	18.8 (29)	0.86 (31)
Mean			-	-	30.7	22	0.99
Range	-	-	-	-	27.2–33.4	18.8–25.0	0.86–1.08
interferon alfa-2b^c	3 MIU TIW	14.4 (30)	8 (27)	134 (31)	4.28 (24) ^d	231.2 (22) ^d	1.40 (35) ^d

PegIFNα2b = peginterferon alfa-2b

a: n = 6, except

b: n = 2

c: n = 16, except

d: n = 6

As shown, t_{1/2} for PegIFNα2b is significantly longer than interferon alfa-2b (30.7 hours versus 4.28 hours respectively). The significantly lower clearance of PegIFNα2b (22.0 mL/hr×kg) relative to interferon alfa-2b (231.2 mL/hr×kg) is reflective of the longer t_{1/2} for the former compound.

Both PegIFNα2b and interferon alfa-2b are rapidly absorbed following subcutaneous administration with mean absorption half-lives (t_{1/2k_a}) of 4.6 hours and 2.3 hours respectively. However, due to sustained maximal concentrations, PegIFNα2b T_{max} is later (range 15-44 hours) compared with interferon alfa-2b (mean 8 hours).

Following absorption, PegIFNα2b displays sustained maximal serum concentrations for 48 to 72 hours post-dose. In contrast, mean serum interferon alfa-2b concentrations decline rapidly after reaching peak concentrations.

Peginterferon alfa-2b C_{max} and AUC increase in a dose-related manner. Mean apparent volume of distribution is slightly higher for interferon alfa-2b (1.4 L/kg) than for PegIFNα2b (0.99 L/kg), which is not anticipated to be of clinical significance.

Upon multiple dosing, there is accumulation of immunoreactive interferons. There is however, only a modest increase in biologic activity as measured by bioassay.

The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. Renal clearance appears to be an important route of elimination for interferon alfa-2b, accounting for approximately 80% of apparent clearance. Based on a retrospective regression analysis of PegIFNα2b Cl/F and creatinine clearance, from an expanded database, it is estimated that renal clearance of PegIFNα2b may account for approximately 30% of the apparent clearance.

Ribavirin plus peginterferon alfa-2b

No PK interactions were noted between PegIFNα2b and RBV in a multiple-dose PK study.

MICROBIOLOGY

Antiviral Activity in Cell Culture

The antiviral activity of BOC was evaluated in a biochemical assay for slow binding inhibitors of NS3/4A protease and in the HCV replicon system. The IC₅₀ and IC₉₀ values for BOC were approximately 200 nM and 400 nM, respectively, in a 72-hour cell culture assay. Loss of replicon RNA appears to be first-order with respect to time of treatment. Treatment at IC₉₀ for 72 hours resulted in a 1-log drop in replicon RNA. Prolonged exposure resulted in a 2-log decrease in RNA levels by Day 15.

Boceprevir cell culture anti-HCV activity was approximately 2-fold lower for an HCV replicon derived from a single genotype 1a isolate, relative to the 1b isolate-derived replicon. In replicon assays, BOC had approximately 2-fold reduced activity against a genotype 2a isolate relative to genotype 1a and 1b replicon isolates. In a biochemical assay, BOC had approximately 3- and 2-fold reduced activity against NS3/4A proteases derived from single isolates representative of HCV genotypes 2 and 3a, respectively, relative to a genotype 1b-derived NS3/4A protease. The presence of 50% human serum reduced the cell culture anti-HCV activity of BOC by approximately 3-fold.

Evaluation of varying combinations of BOC and interferon alfa-2b that produced 90% suppression of replicon RNA showed additivity of effect; no evidence of synergy or antagonism was detected.

Resistance

***In vitro* Studies**

The activity of BOC against the HCV NS3/4A protease or genotype 1b replicon was reduced (2- to 10- fold) by the following amino acid substitutions in the NS3/4A protease domain: V36A/I/M, Q41R, F43C/S, T54A/S, V55A/I, R155K/M/Q, V158I, V170A/T and M175L. A greater than 15-fold reduction in BOC anti-HCV activity was conferred by the substitutions T54C, R155G/I/T and A156S/T/V. The fold decrease in BOC anti-HCV activity conferred by double resistance-associated substitutions was approximately equal to the product of that for the individual substitutions.

Clinical Virology Studies

A pooled analysis was conducted to explore the association between the detection of baseline NS3/4A amino acid polymorphisms and treatment outcome in the two Phase 3 studies, SPRINT-2 and RESPOND-2. Resistance associated polymorphisms were detected in viruses from 6.7% of subjects at baseline; 5.4% had genotype 1a virus and 1.3% had genotype 1b viruses. Overall, the presence of baseline RAVs alone did not appear to have a notable association with treatment response in patients who received the combination of BOC with PegIFN α 2b/RBV.

In a pooled analysis of patients who are previously untreated and patients who have failed previous therapy who received four weeks of PegIFN α 2b/RBV followed by BOC 800 mg TID in combination with PegIFN α 2b/RBV in two Phase 3 studies, post-baseline RAVs were detected in 53% of non-SVR patients. In patients treated with BOC, interferon responsiveness (as defined by

$\geq 1\text{-log}_{10}$ decline in viral load at TW 4) was associated with detection of fewer RAVs, with 6% of these patients having RAVs compared to 41% of patients with $< 1\text{-log}_{10}$ decline in viral load at TW 4 (poorly interferon responsive). In patients treated with BOC with post-baseline samples analyzed for RAVs, interferon responsiveness was associated with detection of fewer RAVs, with 31% of these patients having post-baseline RAVs compared to 68% of patients with $< 1\text{-log}_{10}$ decline in viral load at TW 4. There was no significant difference in the number of non-SVR patients with RAVs detected that received response guided therapy, compared to those that received 48 weeks of BOC/PegIFN α 2b/RBV. The RAVs most frequently detected post-baseline ($> 25\%$ of subjects) in non-SVR subjects were amino acid substitutions V36M (61%) and R155K (68%) in subjects with genotype 1a viruses and T54A (42%), T54S (37%), A156S (26%) and V170A (32%) in subjects with genotype 1b viruses.

Persistence of Resistance-Associated Mutations

Data from an ongoing, long-term follow-up study of subjects who did not achieve SVR in Phase 2 trials with BOC, with a median duration of follow-up of approximately 2 years, indicate that HCV populations harbouring certain post-baseline, BOC-treatment-emergent substitutions may decline in relative abundance over time. However, among those subjects with available data, one or more BOC-treatment-emergent substitutions remained detectable with a population-based sequencing assay in 25% of subjects after 2.5 years of follow-up. The most common NS3/4A substitutions detected after 2.5 years of follow-up were T54S and R155K. The lack of detection of a substitution based on a population-based assay does not necessarily indicate that viral populations carrying that substitution have declined to a background level that may have existed prior to treatment. The long-term clinical impact of the emergence or persistence of BOC-resistance-associated substitutions is unknown. No data are available regarding the efficacy of BOC among subjects who were previously exposed to BOC, or who previously failed treatment with a BOC-containing regimen. The majority of the patients in the long-term follow-up study were not dosed with the indicated BOC regimen.

Effect of Baseline HCV Polymorphisms on Treatment Response

A pooled analysis was conducted to explore the association between the detection of baseline NS3/4A amino acid polymorphisms and treatment outcome in the two Phase 3 studies, SPRINT-2 and RESPOND-2.

Baseline resistance associated polymorphisms were detected in 7% of subjects by a population-based sequencing method. Overall, the presence of these polymorphisms alone did not impact SVR rates in subjects treated with BOC. However, among subjects with a relatively poor response to PegIFN α 2b/RBV during the 4-week lead-in period, the efficacy of BOC appeared to be reduced for those who had V36M, T54A, T54S, V55A or R155K detected at baseline. Subjects with these baseline polymorphisms and reduced response to PegIFN α 2b/RBV represented approximately 1% of the total number of subjects treated with BOC.

Cross-Resistance

Many of the treatment-emergent NS3/4A amino acid substitutions detected in BOC-treated subjects who did not achieve SVR in the Phase 3 clinical trials have been demonstrated to reduce

the anti-HCV activity of other HCV NS3/4A PIs. The impact of prior exposure to BOC or treatment failure on the efficacy of other HCV NS3/4A PIs has not been studied. The efficacy of BOC has not been established for patients with a history of exposure to other NS3/4A PIs. Cross-resistance is not expected between BOC and interferons, or BOC and RBV.

Pharmacogenomics

Genetic variance near the gene encoding interferon-lambda-3 (*IL28B rs12979860*, C to T change) has been demonstrated to be a strong predictor of response to PegIFN α 2b/RBV. *IL28B rs12979860* was genotyped in 653 of 1048 (62%) subjects in SPRINT-2 (previously untreated) and 259 of 394 (66%) subjects in RESPOND-2 (previous treatment failure) (see **CLINICAL TRIALS**).

Overall, among the patients that received at least one dose of placebo or BOC (Modified-Intent-to-Treat population) plus PegIFN α 2b/RBV for 48-weeks, the SVR rates tended to be lower in patients with the C/T and T/T genotypes when compared to the C/C genotype of both Previously Untreated Patients and Previous Treatment Failures. The results of this retrospective subgroup analysis should be viewed with caution because of the small sample size and potential differences in demographic or clinical characteristics of the sub-study population relative to the overall trial population.

Table 29: Sustained Virologic Response Rates by *IL28B rs12979860* Genotype

Clinical Study	<i>IL28B rs12979860</i> Genotype	SVR, % (n/N)		
		PegIFN α 2b/RBV-48 ^a	BOC/PegIFN α 2b/RBV (RGT)*	BOC/PegIFN α 2b/RBV-48 ^a
SPRINT-2 (Previously Untreated Subjects)	C/C	78 (50/64)	82 (63/77)	80 (44/55)
	C/T	28 (33/116)	65 (67/103)	71 (82/115)
	T/T	27 (10/37)	55 (23/42)	59 (26/44)
RESPOND-2 (Subjects Who Have Failed Previous Therapy)	C/C	46 (6/13)	79 (22/28)	77 (17/22)
	C/T	17 (5/29)	61 (38/62)	73 (48/66)
	T/T	50 (5/10)	55 (6/11)	72 (13/18)

a: For description of each treatment arm, see **CLINICAL TRIALS**.

TOXICOLOGY

Acute Toxicity

Boceprevir

Boceprevir has a low order of oral single-dose toxicity. No adverse effects were observed in rats at doses of 2,000 mg/kg in females and 1,000 mg/kg in males. In monkeys, no adverse effects were observed at doses up to 250 mg/kg and only minor clinical signs of emesis and abnormal stool findings were observed at 500 and 1,000 mg/kg. No adverse effects were observed in dogs at a dose of 100 mg/kg and the only findings observed at the highest dose tested of 200 mg/kg were emesis and abnormal stool findings.

Ribavirin

Ribavirin has a low level of toxicity after single doses in animals. Depression, hunched posture, rough coat, ataxia, prostration, tremors and labored respiration were observed following acute toxic doses in mice and rats. The presence of dark red material in the gastro-intestinal tract of non-surviving rats in the higher dose groups is suggestive of RBV -related hemorrhage. Similar drug-related findings were observed in guinea pigs. Soft, mucoid stools and/or diarrhea were observed in dogs administered single doses up to 480 mg/kg. Emesis and transient body weight loss were seen in select dogs at the intermediate doses. Ulcer-like areas in the upper portion of the duodenum were observed in dogs treated with the 60/480 mg/kg dose.

Peginterferon alfa-2b

Volume limited single subcutaneous doses of PegIFN α 2b (56,850 mcg/m² in rats and 60,410 mcg/m² in mice) failed to produce any toxic effects. Single dose studies in primates showed a no-observed adverse effect level of 58,861 mcg/m² for males and 29,435 mcg/m² for females. These single dose no-observed adverse event levels correspond with approximately 2,943 (for male monkeys) and 1,472 (for female monkeys) times the weekly human dose of interferon alfa-2b (20 mcg/m² or 9 MIU) utilized in the treatment of chronic hepatitis C. Two of four monkeys given a single dose of 117,721 mcg/m² either died or was sacrificed in moribund condition. One death was the result of myocarditis and pleural effusion that was most likely present prior to dosing. The cause of death in the other monkey was not determined. Adverse events noted in survivors at the high-dose included loss of appetite, lower body temperature and reduction of blood pressure.

Long-term (subacute and chronic) Toxicity

Boceprevir

Boceprevir-related nonclinical toxicology findings tend to be dose-related and/or species-specific in nature; and occur at exposures that are slightly higher than or similar to the clinical therapeutic dose. Targets identified in nonclinical studies consist of the gall bladder, liver, reproductive tract, and the activated partial thromboplastin time (APTT) coagulation parameter; and were primarily observed in rodents. None of these nonclinical findings have been identified clinically.

Discolored gall bladder was observed in the 2-year mouse carcinogenicity study with no histopathologic correlate, inflammation, evidence of concretions, long-term impact to gall bladder integrity or gall bladder tumors. In shorter duration mouse studies (at higher doses and exposures) and in monkey studies, no gall bladder discoloration was observed. The NEL for this finding represents a mouse-to-human exposure multiple of less than 1-fold the human exposure at the human therapeutic dose of 800 mg TID. There have been no BOC-related clinical AEs suggestive of impaired gall bladder function.

Nonclinical toxicology findings related to the liver consisted of focal neutrophilic infiltrates (mice), single cell hepatocyte necrosis (mice), pigment accumulation in liver macrophages (mice), and multinucleated hepatocytes (male rats). With the exception of the multinucleated hepatocytes, all liver-related finding NELs represent mouse-to-human or rat-to-human exposure multiples similar to or greater than the human exposure at the human therapeutic dose of 800 mg TID. Liver function tests have revealed no evidence of clinical concern.

Increases in the coagulation parameter APTT have been observed in monkeys. There were no changes in other clinical pathology parameters that would be suggestive of hemorrhage, or gross pathology evidence of hemorrhage that would indicate a defect in hemostasis. The NEL for this finding represents a monkey-to-human exposure multiple of less than 1-fold the human exposure at the human therapeutic dose of 800 mg TID. APTT levels have been monitored in the clinic, and no clinically meaningful effects have been identified.

Ribavirin

Repeated dose toxicity studies were conducted in mice, rats, dogs and monkeys. In mice administered oral doses up to 600 mg/kg for 90 days, dose-related mortality was observed in the highest dose groups (300 and 600 mg/kg/day). Dose-related decreases in hemoglobin, hematocrit and/or erythrocyte count were observed in all treated groups. Increases in mean corpuscular volume/hemoglobin (300 mg/kg), platelets (150, 300 mg/kg/day), reticulocyte values (300 mg/kg/day), neutrophils, nucleated RBCs and poikilocytes (300 mg/kg/day) were also observed. Treated animals had higher spleen weights and increased liver weight was observed in the 300 mg/kg group. Testes and epididymides of mice in the 150, 300 and 600 mg/kg dose groups had histological evidence of bilateral degeneration of the testicular germinal epithelium accompanied by bilateral epididymal oligospermia and accumulation of degenerated seminal product.

No RBV -related mortality or clinical observations were observed in rats administered oral RBV in doses up to 120 mg/kg for at least 28 days. Food consumption and body weights were decreased in the mid to high dose groups. A treatment-related decrease in circulating red cell mass characterized by decreased hematocrit, hemoglobin and erythrocyte counts was observed. Small thymus and lymphoid depletion in the thymus of high dose rats of both sexes was also observed.

In a ninety-day study in rats administered oral RBV (20 to 200 mg/kg/day); mortalities were observed in the 15 mg/kg/day (1F) and 200 mg/kg/day (1 F/3 M) dose groups. Increased circulating aspartate aminotransferase and alanine aminotransferase were observed in the highest dose group in addition to the treatment-related effects on hemoglobin, hematocrit, erythrocyte count, platelet and leukocyte count (40 - 200 mg/kg). Heart and spleen weights were elevated in rats in the highest dose group.

In a 30-day dietary study in rats who received RBV up to 320 mg/kg/day, piloerection, abnormal breathing and/or subdued behavior was observed in animals that died or were sacrificed prematurely. Hunched posture, pale extremities, ataxia, thin build, hypothermia, body weight loss or reduced weight gain and reduced food consumption were observed in the 160 and 320 mg/kg dose groups. Hematological effects included reduced hemoglobin and hematocrit (40 and 160 mg/kg), reduced RBC and WBC counts (160 mg/kg), decreased mean corpuscular hemoglobin and mean corpuscular volume (10 and 40 mg/kg). Females in the 160 mg/kg dose group showed increases in AST and phosphate and decrease in total protein, albumin, creatine and calcium. Decreases in thymus weight and increases in heart and lung weights occurred at 160 mg/kg. Macroscopic changes occurred in the heart, liver, lungs, thorax, thymus and lymph nodes (160 mg/kg) with histological changes in the intestine, heart, liver, lungs, salivary glands, spleen, testes, thymus and lymph nodes. Similar findings were noted in a 52-week dietary study where rats were administered up to 90 mg/kg/day. In addition to the findings noted above, ALT levels were reduced (both sexes) and a reduction in albumin noted in females. Albumin was increased in males (90 mg/kg) between weeks 4 and 13 and reduced in weeks 26 to 52. Pathology findings revealed depletion in thymic cellularity with possible secondary findings in the skin, lung and submandibular lymph nodes.

In a 28-day study in dogs administered oral RBV doses of 15, 30 or 60 mg/kg/day, 5 of 6 dogs in the high dose group died or were sacrificed due to moribund condition. Enteritis was the principal effect noted in the low dose group whereas body weight and food consumption were decreased primarily in the mid and high dose groups. Severe cachexia was noted in the high dose group. There was evidence of non-regenerative anemia (decreased red cell mass and absence of reticulocytosis) and decreased white blood cells in the mid- and high-dose groups. Bone marrow hypoplasia, lymphoid depletion in the thymus, gastric cytoplasmic vacuolization, and increased brown granular pigment in the spleen were present in mid- and high-dose groups. Similar findings were noted in two additional studies where dogs were administered oral doses of 5- 40 mg/kg/day for 30 days or 5 - 20 mg/kg/day for 52 weeks.

Ribavirin was well tolerated when administered to cynomolgus monkeys at a single daily oral (gavage) dose of 100 mg/kg for two weeks. Upon intramuscular administration (30 or 100 mg/kg) to rhesus monkeys for 10 days, reversible, dose-related decreases in circulating red blood cell mass was observed as well as an increase in megakaryocytes at day 10 in both groups.

Peginterferon alfa-2b

In a multiple dose study, doses of 1,414 mcg/m² and 4,239 mcg/m² administered every other day over one month to primates were also well tolerated. At the highest dose (14,126 mcg/m² QOD) one female was sacrificed in moribund condition with abscesses at the injection site due to bacterial infection, bone marrow hypocellularity, anemia, hypoalbuminemia and lymphoid depletion in thymus, spleen and lymph nodes. In surviving monkeys, there were dose related decreases in platelets, neutrophils, lymphocytes, erythrocytes, serum proteins, calcium, phosphorus and potassium. These changes tended toward normal after Week 2, and returned to pre-test values in the high-dose monkeys retained for a 4-week recovery period. Test article-related histopathologic findings were restricted to the bone marrow and injection site. There was minimal to mild bone marrow hypercellularity, suggestive of a regenerative process, in some animals in the PegIFN α 2b and the interferon alfa-2b groups, and minimal to mild perivascularitis at injection sites in all groups, including vehicle controls. All of the changes that were seen in PegIFN α 2b-dosed monkeys were similar in nature to those seen in the interferon alfa-2b group of this study, or in previous studies in monkeys of interferon alfa-2b.

The table below (Table 30) illustrates the relationships between dose, general effects, and AUC for interferon alfa-2b after a single dose in the one-month study in monkeys.

The general toxicologic effects of a dose of 4,239 mcg/m² of PegIFN α 2b are most similar to those of an interferon alfa-2b dose of 3,105 mcg/m². Peginterferon alfa-2b was well tolerated but produced mild toxicity at the mid-dose of 4,239 mcg/m², or 241.6 MIU/m². Interferon alfa-2b was well tolerated but produced similar mild toxicity at a dose of 3,105 g/m², or 807.3 MIU/m². These doses gave similar interferon alfa-2b plasma levels in terms of AUC (0.675 vs. 0.545 MIU-equiv.x hr/mL) after a single dose. However, PegIFN α 2b accumulates after multiple doses because of its long half-life. The plasma AUC for a PegIFN α 2b dose of 4,239 mcg/m² would exceed that of an interferon alfa-2b dose of 3,105 mcg/m² after multiple doses. This suggests that the severity of general toxicologic effects in monkeys may be somewhat less for PegIFN α 2b than for interferon alfa-2b at similar systemic exposure levels.

Table 30: Correlation of Dose and General Effects

Test Article	Dose (mcg/m ²)	MIU/m ²	IFN AUC/(I) ^c MIU-equiv. x hr/mL (after one dose)	Effect
PegIFN α 2b ^a	1,414	80.6	0.284	Well tolerated, mild toxicity
PegIFN α 2b	4239	241.6	0.675	Well tolerated, minimal effects
PegIFN α 2b	14,126	805.2	2.22	Well tolerated, mild toxicity, moderate weight loss
interferon alfa-2b ^b	3,105	807.3	0.545	One female sacrificed moribund Moderate effects in other animals

PegIFN α 2b = peginterferon alfa-2b

a: Based on a specific activity of 0.57 X 10⁸ IU/mg protein

b: Based on a specific activity of 2.6 X 10⁸ IU/mg protein

c: As measured by ELISA

Peginterferon alfa-2b plus ribavirin

A one-month toxicity study and a one-month toxicity and neutrophil function study investigating PegIFN α 2b in combination with RBV were conducted in cynomolgus monkeys. In each study, monkeys were dosed with subcutaneous injections of PegIFN α 2b every other day (5,494 mcg/m²) either alone or in combination with a daily oral (gavage) RBV (50 or 75 mg/kg).

The major effects observed in the one-month toxicity study (body weight loss related to inappetence, and decreased red and white blood cell parameters due to reversible bone marrow suppression) were those that were expected based on the known biologic effects of these two drugs. Five monkeys (one peginterferon monotherapy, one low-dose combination, and three high-dose combination) were sacrificed in moribund condition. The moribundity observed was not a direct test article-related effect but was due to opportunistic infections. Peginterferon alfa-2b's effect on neutrophil counts supports the conclusion that lowered host resistance to infections was a contributing factor in the morbidity and mortality. In the follow-up study, a similar degree of morbidity due to opportunistic infections was not observed with similar combination dosing regimens.

In the one-month toxicity and neutrophil function study, administration of RBV (75 mg/kg) in combination with PegIFN α 2b (350, 1,400, or 5,500 mcg/m²) resulted in an exacerbation of biologic effects typically seen with either drug alone. These findings included body weight loss related to inappetence, and decreased red and white blood cell parameters due to reversible bone marrow suppression (PegIFN α 2b) or extravascular hemolysis (RBV). Bone marrow changes were noted for all three combination groups with the highest incidence occurring in the high-dose combination males. There were no drug-induced effects on neutrophil function. One monkey died of aspiration pneumonia most likely exacerbated by neutropenia.

Carcinogenicity and Mutagenicity

Boceprevir

Two-year carcinogenicity studies in mice and rats were conducted with BOC. Mice were administered doses up to 650 mg/kg. Rats were administered doses of up to 125 mg/kg in males and 100 mg/kg in females. At the high dose of 650 mg/kg in female mice, the incidence of hepatocellular adenomas was increased at systemic exposures 5.7-fold of those in humans at the recommended 800 mg TID clinical dose. There were no increases in mortality or malignancy associated with the hepatocellular adenomas. Induction of CYP450 enzymes has been demonstrated previously in mice administered BOC, and liver tumours are a recognized sequelae with chronic exposure to an enzyme inducer. There were no increases in the incidence of tumours in male mice at any dose in the study. In rats, no treatment-related increase in adenomas or carcinomas occurred at systemic exposures similar to the human exposure at the recommended 800 mg TID clinical dose (rat-to-human multiple of ~ 0.9-fold). The clinical relevance of the hepatocellular adenomas observed in female mice, if any, is unknown.

Boceprevir was not mutagenic or genotoxic in a battery of *in vitro* or *in vivo* assays, including bacterial mutagenicity, human peripheral blood lymphocyte and mouse micronucleus assays.

Ribavirin

An 18-month study in mice indicated that RBV was not oncogenic at oral gavage doses up to 75 mg/kg/day. Hematologic and splenic changes indicate that RBV was absorbed at sufficient levels to cause adverse effects, and that the severity of the effects was dose-related. A two-year study in rats indicated that RBV had no oncogenic effect at oral gavage doses up to 40 mg/kg/day. Conventional carcinogenicity studies in rodents with low exposures compared to human exposure under therapeutic conditions (factor 0.1 in rats and 1 in mice) did not reveal tumorigenicity of RBV. In addition, in a 26 week carcinogenicity study using the heterozygous p53 (+/-) mouse model, RBV did not produce tumors at the maximally tolerated dose of 300 mg/kg (plasma exposure factor approximately 2.5 compared to human exposure).

Mutagenicity studies suggest that RBV may exert some mutagenic activity. Ribavirin induced an increase in transformed foci in two independent Balb/3T3 in vitro transformation assay trials. Mutagenic activity was observed in the mouse lymphoma forward mutation assay, and RBV induced micronuclei in polychromatic erythrocytes at doses of 20-200 mg/kg in a mouse micronucleus assay. A dominant lethal assay in rats was negative, indicating that if mutations occurred in rats they were not transmitted through male gametes.

Peginterferon alfa-2b

Studies with PegIFN α 2b have not been performed to determine carcinogenicity due to lack of biologic activity in rodents.

Mutagenicity studies with PegIFN α 2b revealed no mutagenic activity.

Reproductive and Developmental Toxicity

Boceprevir

Following a single, oral dose of 30 mg/kg ¹⁴C-BOC, medicine-derived radiocarbon was transferred into the milk of lactating, 12-day postpartum rats. Peak systemic concentrations of medicine-derived radiocarbon in nursing pups were over 100-fold lower than in the mothers. Exposure to medicine-derived materials in nursing human infants is estimated to be less than 1% of the dose.

In rats, BOC induced reversible effects on fertility and early embryonic development in female rats with a No Effect Level (NEL) of 75 mg/kg. At this dose, the rat-to-human exposure multiple is 1.3-fold higher than the systemic human exposure at the recommended human therapeutic dose of 800 mg TID. Decreased fertility was also observed in male rats, most likely as a consequence of testicular degeneration (included Sertoli cell vacuolation, atrophy of the seminiferous tubule epithelium, epididymal cellular debris, hypospermia of the epididymides, spermatid degeneration, soft testes, and decreased sperm count and motility), with evidence of partial reversibility and a NEL of 15 mg/kg (which represents a rat-to-human exposure multiple of less than 1-fold the human exposure at the human therapeutic dose of 800 mg TID). In juvenile rats, the mean age of attainment of balanopreputial separation was minimally higher compared to the control group (46.1 compared to 44.4 days). This finding was also associated with testicular/epididymal findings similar to those observed in adult rats, and included lower mean epididymides and testes weights.

The NEL for these findings represents a rat-to-human exposure multiple of less than 1-fold the human exposure at the human therapeutic dose of 800 mg TID. Testicular degeneration has not been observed in mice or monkeys and therefore is considered species-specific to rats. Additionally, clinical monitoring of the surrogate marker inhibin-B, as well as semen analysis has revealed no evidence that these findings are clinically relevant in human patients.

In juvenile rats, BOC induced reversible hyperplasia of thyroid gland follicular cells and higher liver weights, with a NEL that represents a rat-to-human exposure multiple of less than 1-fold the human exposure at the human therapeutic dose of 800 mg TID. Since no BOC-related thyroid hyperplasia has been observed in any adult rat, mouse or monkey studies, these changes are considered age- and species-specific to the juvenile rat, and not a safety concern for humans. Thyroid function tests have been included in BOC clinical studies with no evidence of BOC-related thyroid changes.

Ribavirin

In a Segment I study with male rats administered 10, 40, or 160 mg/kg daily for 9 weeks prior to mating and female rats administered 0.3, 1.0 or 10.0 mg/kg daily (all doses lower than the recommended 24-hour human dose for RBV) from 2 weeks prior to mating until Day 5 of presumed pregnancy, there were no treatment-related effects on female mating performance or latent effects on *in utero* and postnatal development of the litter.

Rats were administered oral doses of 0.3, 1.0 and 10.0 mg/kg RBV from day 6 to day 15 of pregnancy. Despite a negligible maternal toxicity response, the 10.0 mg/kg dose resulted in embryonic death, reduced litter size, reduced fetal weight, visceral and skeletal abnormalities, and skeletal variants. Morphological changes were also observed in the 1.0 mg/kg dose group. In a second study, rats were administered oral doses of 0.1, 0.3 and 1.0 mg/kg RBV from day 6 to day 18 of pregnancy. The 1.0 mg/kg dose level was associated with a transient reduction in maternal weight gain, slight reduction in fetal weight and increased incidences of fetuses showing skeletal anomalies and variants.

Male mice were administered oral RBV (0, 35, 75, 150 mg/kg/day) for 3-6 months to evaluate the time course and reversibility of RBV -induced testicular degeneration. Moderate dose-dependent effects on spermatogenesis were observed including reduced testicular spermatid counts, reduced sperm motility, increased abnormal epididymal sperm, reduced testes weight, marginal reductions in seminiferous tubular diameter and increased incidences of vacuolization and reduced thickness of the germinal cell epithelium. Upon cessation of treatment, total recovery was observed within one to two spermatogenesis cycles. In a second study, RBV administered at daily oral doses of 1, 15, 35 or 75 mg/kg for 3 months to seven-week old mice resulted in dose-related decreases in testicular spermatid and caudal epididymal sperm counts and in the percentage of morphologically normal sperm. Morphological changes were the most sensitive endpoint observed with differences from control detected at 15 mg/kg RBV. The effects on spermatid count and sperm morphology were reversible upon cessation of dosing.

A toxicokinetic study was conducted in rabbits at oral doses of 0.3, 1.0 and 3.0 mg/kg from day 6 to day 18 of pregnancy. Effects on the developing conceptus included, at 3.0 mg/kg, a slight increase in *in utero* mortality, slight reduction in litter size, markedly lower litter and mean fetal weight and an increase in fetuses showing probable malformation upon gross examination. Rats administered RBV at doses up to 1.0 mg/kg from day 15 of pregnancy through day 21 postpartum produced little or no evidence of toxicity to either the pregnant female or her offspring.

Peginterferon alfa-2b

Interferon alfa-2b has been shown to have abortifacient effects in *Macaca mulatta* (rhesus monkeys) at 7.5, 15, and 30 million IU/kg (90, 180, and 360 times the intramuscular or subcutaneous dose of 2 million IU/m²). Although abortion was observed in all dose groups, it was only statistically significant at the mid- and high-dose groups. Peginterferon alfa-2b is likely to also cause these effects.

REFERENCES

1. Lindsay KL, Trepo C, Heintges T, Shiffman ML, Gordon SC, Hoefs JC, et al. A Randomized, Double-Blind Trial Comparing Pegylated Interferon Alfa-2b to Interferon Alfa-2b as Initial Treatment for Chronic Hepatitis C. *Hepatology* 2001; 34 (2):395-403.
2. Hoffmann RM, Diepolder HM, Zachoval R, Zwiebel F-M, Jung MC, Scholz S, et al. Mapping of immunodominant CD4+ T lymphocyte epitopes of hepatitis C virus antigens and their relevance during the course of chronic infection. *Hepatology* 1995;21(3):632-8.
3. Knodell RG, Ishak KG, Black WC, Chen TS, Craig R, Kaplowitz N, et al. Formulation and application of a numerical scoring system for assessing histological activity in asymptomatic chronic active hepatitis. *Hepatology* 1981;1(5): 431-5.
4. Vial T, Descotes J. Clinical toxicity of the interferons. *Drug Saf* 1994;10(2):115-50.
5. National Institutes of Health Consensus Development Conference Panel Statement: Management of Hepatitis C. *Hepatology* 1997;26(3 Suppl. 1):2-10.
6. Center for Disease Control (CDC): Recommendations for Prevention and Control of Hepatitis C Virus (HCV) Infection and HCV-Related Chronic Disease. *MMWR Recomm* 1998; 47(RR19): 1-39.
7. Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001; 358(9286):958-65.
8. McHutchison JG, Manns M, Patel K, Poynard T, Lindsay KL, Trepo C, et al. Adherence to Combination Therapy Enhances Sustained Response in Genotype-1-Infected Patients with Chronic Hepatitis C. *Gastroenterology* 2002; 123(4):1061-9.

9. Gupta SK, Glue P, Jacobs S, Belle D, Affrime M. Single-dose pharmacokinetics and tolerability of pegylated interferon-alpha 2b in young and elderly healthy subjects. *Br J Clin Pharmacol* 2003; 56(1):131-4.
10. M. Sherman, S. Shafran, K. Burak et al. Management of chronic hepatitis C: Consensus guidelines. *Can. J. Gastroenterol.* 2007; 21 (Suppl C): 25C - 34C.
11. Chase R., Skelton A., Xia E., Curry S., Liu S., McMonagle P., et al. A novel HCV NS3 protease mutation selected by combination treatment of the protease inhibitor boceprevir and NS5B polymerase inhibitors. *Antiviral Res.* 84(2): 178-84, 2009.
12. Harrison SA, Rossaro L., Hu KQ, Patel K., Tillmann H., Dhaliwal S., et al. Serum cholesterol and statin use predict virological response to peginterferon and ribavirin therapy. *Hepatology.* 52 (3): 864-74, 2010.
13. Kwo P., Lawitz E., McCone J., Schiff E., Vierling J., Pound D., et al. Efficacy of Boceprevir, an NS3 Protease Inhibitor, in Combination with Peginterferon alfa-2b and Ribavirin in Treatment-Naive Patients with Genotype 1 Hepatitis C Infection (SPRINT-1): an Open-Label, Randomised, Multicentre Phase 2 Trial. *Lancet.* 376:705-16, 2010.
14. Qiu P., Sanfiorenzo V., Curry S., Guo Z., Liu S., Skelton A. et al. Identification of HCV protease inhibitor resistance mutations by selection pressure-based method. *Nucleic Acids Res.* 37 (10): e74, 2009.
15. Susser S., Welsch C., Wang Y., Zettler M., Domingues FS, Karey U., et al. Characterization of resistance to the protease inhibitor boceprevir in hepatitis C virus-infected patients. *Hepatology.* 50:1709-18, 2009.
16. Poordad F., McCone J., Bacon B., Bruno S., Manns M., Sulkowski M., et al. Boceprevir for Untreated Chronic HCV Genotype 1 Infection. *N Engl J Med.* 2011; 364:1195-1206.
17. Bacon B., Gordon S., Lawitz E., Marcellin, P., Vierling J., Zeuzem S., et al. Boceprevir for Previously Treated Chronic HCV Genotype 1 Infection. *N Engl J Med.* 2011; 364:1207-17.
18. PEGETRON[®] (ribavirin + peginterferon alfa-2b) Product Monograph, March 22, 2011.
19. VICTRELIS[™] (boceprevir) Product Monograph, July 27, 2011.

PART III: CONSUMER INFORMATION

VICTRELIS TRIPLE™

VICTRELIS™
boceprevir

PLUS

PEGETRON®
ribavirin
plus
peginterferon alfa-2b

This leaflet is part III of a three-part “Product Monograph” published when VICTRELIS TRIPLE™ was approved for sale in Canada and is designed specifically for Consumers. Please read this information carefully before starting your VICTRELIS TRIPLE™ therapy. It is important to read this each time your prescription is refilled in case new information becomes available. This leaflet is a summary and will not tell you everything about VICTRELIS TRIPLE™. Contact your doctor or pharmacist if you have any questions about the drug.

You should talk with him or her before starting therapy and at your regular check-ups. This medicine has been prescribed for you personally and you should not pass it on to others. It may harm them, even if their symptoms are the same as yours.

ABOUT THIS MEDICATION

What the medication is used for:

VICTRELIS TRIPLE™ is a prescription medicine used to treat chronic (lasting a long time) hepatitis C genotype 1 infection in adults who have not been previously treated or who have failed previous therapy. Patients with hepatitis C have the virus in their blood and in their liver.

It is not known if VICTRELIS TRIPLE™ is safe and effective when used in children (< 18 years of age).

What it does:

VICTRELIS TRIPLE™ therapy consists of 3 active medications in 2 drug products: VICTRELIS™ (boceprevir) and PEGETRON®, (ribavirin and peginterferon alfa-2b).

Boceprevir is a medicine called a Hepatitis C Virus Protease Inhibitor that directly targets Hepatitis C Virus to reduce the amount of virus in your body.

Ribavirin is an antiviral agent (fights infection), but does not work when used by itself to treat chronic hepatitis C.

Peginterferon alfa-2b generally helps the body’s immune system to fight infections.

It is not known exactly how ribavirin and peginterferon alfa-2b work together to fight the hepatitis C infection.

VICTRELIS TRIPLE™ therapy may reduce the amount of hepatitis C virus in the blood stream to below the level that can be measured by a laboratory test.

When it should not be used:

Do not use VICTRELIS TRIPLE™:

- **If you or your partner are pregnant.**
- **If you or your partner plan to become pregnant during treatment or during the 6 months after treatment.**
- **If you or your partner become pregnant during treatment.** VICTRELIS TRIPLE™ therapy can cause serious birth defects or harm to your unborn child. Therefore, both you and your partner **must use effective contraception** during this time.
- If you are allergic to any of the ingredients in VICTRELIS TRIPLE™ (boceprevir, ribavirin or peginterferon alfa-2b) or to any of the excipients (see **What the non-medicinal ingredients are**).
- If you have autoimmune hepatitis (hepatitis caused by cells in your body attacking each other) or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have a severe nervous or mental disorder.
- If you have thyroid disease that is not well controlled with medicines.
- If you have advanced uncontrolled liver disease (other than hepatitis C).
- If you have severe kidney disease.
- If you have epilepsy.
- If you are breastfeeding.
- If you are taking certain medicines. For more information about medicines that you should not take while using VICTRELIS TRIPLE™, please see **INTERACTIONS WITH THIS MEDICATION**.

What the medicinal ingredients are:

- boceprevir capsules
- ribavirin capsules
- peginterferon alfa-2b powder for solution

What the non-medicinal ingredients are:

Boceprevir capsules: Non-medicinal ingredients are croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline

cellulose, pre-gelatinized starch, and sodium lauryl sulfate. The capsule shell consists of gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. The capsule is printed with red ink. The red ink contains red iron oxide and shellac.

Ribavirin capsules: Non-medicinal ingredients are croscarmellose sodium, lactose monohydrate, magnesium stearate and microcrystalline cellulose. The capsule shell contains gelatin, sodium lauryl sulfate, silicon dioxide, and titanium dioxide.

Peginterferon alfa-2b Powder for Solution in REDIPEN® single dose delivery system: Non-medicinal ingredients are polysorbate 80, sodium phosphate dibasic anhydrous, sodium phosphate monobasic dihydrate, and sucrose.

What dosage forms it comes in:

Each boceprevir capsule contains 200 mg of boceprevir.

Each ribavirin capsule contains 200 mg of ribavirin.

Peginterferon alfa-2b REDIPEN® single dose delivery system consists of a dual-chamber glass cartridge with a chamber containing peginterferon alfa-2b as a white to off-white lyophilized powder and another chamber containing Sterile Water for Injection. The cartridge is provided in a pen device for reconstitution, dose preparation and subcutaneous administration. It is available in strengths of 80, 100, 120, and 150 mcg for single use.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Some people get depressed when taking pegylated interferon alfa-2b alone or in combination treatment with ribavirin, and in some cases people had thoughts about threatening the life of others, suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

What is the most important information I should know about VICTRELIS TRIPLE™ therapy?

1. VICTRELIS TRIPLE™ therapy could cause serious birth defects or harm your unborn child.

- **If you or your partner are pregnant, you should not receive VICTRELIS TRIPLE™.**
- **Pregnancy should not be planned while you or your partner are on therapy or for 6 months after therapy.**
- **If you or your partner become pregnant while on**

therapy or during the 6 months after stopping therapy, consult your doctor immediately. Your doctor should call the Merck Canada Inc. Medical Information Department at 1-800-567-2594.

- **If you are a woman of childbearing age, you must have a negative pregnancy test before treatment and a pregnancy test each month during treatment.**
 - **Both you and your partner must use effective contraception during treatment and for the 6 months after treatment is completed.** As systemic (e.g., oral, topical...) hormonal contraceptives may not work as well while taking VICTRELIS TRIPLE™, use of 2 alternative methods of contraception, such as barrier method and intrauterine devices during treatment with VICTRELIS™ and ribavirin. **You should discuss with your doctor how you or your partner can prevent getting pregnant.**
2. **Ribavirin may cause anemia, which is a decrease in the number of red blood cells you have. Anemia may be increased when boceprevir is added to peginterferon alpha/ribavirin therapy. This can be dangerous, especially for patients who already have heart or circulatory (cardiovascular) problems. Talk with your doctor before taking VICTRELIS TRIPLE™ therapy if you have or have ever had any cardiovascular problems. Your healthcare provider will be checking your blood counts periodically for possible decreases in your blood count. Depending on the medications that you are taking, your healthcare provider may make changes to your current medicines or prescribe additional medicines to treat your anemia.**
 3. **Boceprevir may cause serious side effects when taken with other medications. It is important to know the medicines that should not be taken with boceprevir.**
 4. **Do not take boceprevir alone to treat chronic hepatitis C infection. Boceprevir should only be used with other medicines to treat chronic hepatitis C infection.**

Medicines are sometimes prescribed for purposes other than those listed in this package leaflet. Remember, this medicine is for you and must be used as prescribed by your doctor. Never give it to anyone else.

BEFORE you use VICTRELIS TRIPLE™ talk to your doctor or pharmacist if you have any of the following medical conditions or other serious medical problems:

- Previous heart attack, or other heart problems, because therapy may cause heart problems to become worse.
- Blood disorders; including anemia (low red blood cell count), thalassemia (Mediterranean anemia), sickle-cell anemia, and neutropenia (low neutrophil count) because therapy may further reduce the number of red and white blood cells you have. This

may make you feel dizzy or weak and could worsen any heart problems you might have.

- Kidney problems.
- Liver problems (except hepatitis C infection).
- Nervous or mental problems (such as depression, anxiety, etc.), because the therapy could make these problems worse.
- Body organ transplant and are taking medicine that keeps your body from rejecting your transplant (suppresses your immune system).
- Thyroid disease.
- Cancer.
- Infection with hepatitis B virus and/or Human Immunodeficiency Virus (the virus that causes AIDS).
- If you have had problems with your immune system.
- If you have diabetes or high blood pressure, your doctor may ask you to have periodic eye examinations.
- If you have high blood fat levels (such as elevated triglycerides or cholesterol levels).
- If you had any serious illness affecting your breathing or your blood.
- If you have psoriasis or sarcoidosis, it may become worse while you are using VICTRELIS TRIPLE™.
- If you have any other medical condition.
- Be sure to tell your doctor about all the medications you are taking, including those without a prescription, and the Chinese herbal medication Shosaikoto (also known as Xiao-Chai-Hu-Tang).
- If you are using any other medications.

VICTRELIS TRIPLE™ may cause a Reduction of Red Blood Cells a condition known as anemia, or a reduction of neutrophils (a type of white blood cell) a condition known as neutropenia. Anemia or neutropenia may be increased when VICTRELIS™ is added to your ribavirin therapy. Therefore, your healthcare provider will be checking your blood counts periodically for possible decreases in your blood cell counts. Depending on the medications that you are taking, your healthcare provider may make changes to your current medicines or prescribe additional medicines to treat your anemia, or neutropenia.

VICTRELIS TRIPLE™ may cause serious side effects when taken with other medications. It is important to know the medicines that should not be taken with VICTRELIS™.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients who received VICTRELIS TRIPLE™. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with VICTRELIS TRIPLE™. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

INTERACTIONS WITH THIS MEDICATION

Drugs that may interact with ribavirin and/or peginterferon alfa-2b include: medications metabolized by CYP1A2, CYP2C8/9 and CYP2D6; reverse transcriptase inhibitors such as zidovudine and stavudine; purine nucleoside analogues such as didanosine and abacavir; and Highly Active Anti-Retroviral Therapy.

Co-administration of ribavirin capsules and didanosine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

Tell your doctor or pharmacist if you are taking SEBIVO* (telbivudine) for chronic hepatitis B because taking this medicine together with pegylated interferon alfa-2b may increase your risk of developing peripheral neuropathy (numbness, weakness, tingling, and/or burning sensations, or pain in the arms and/or legs). The combined use of these medications is not recommended.

Do not take VICTRELIS TRIPLE™ if you take:

- alfuzosin – used to treat enlarged prostate;
- amiodarone, propafenone and quinidine – used for heart beat problems;
- astemizole¹, terfenadine¹ – used to treat allergies, hives, itching and watery eyes;
- birth control pills that contain drospirenone;
- carbamazepine, phenobarbital, phenytoin – used to treat seizures and nerve pain;
- cisapride¹ – used to help with digestion;
- ergot-containing medicines used to treat migraines, such as:
 - ergotamine,
 - dihydroergotamine,
 - ergonovine,


¹ Please note that cisapride, astemizole and terfenadine are no longer available on the Canadian market.

- methylergonovine;
- lovastatin, simvastatin – used for lowering high cholesterol and triglycerides;
- oral midazolam, oral triazolam – used to help you sleep;
- pimozide – used for mental health problems;
- rifampin – used to treat tuberculosis or meningitis;
- sildenafil and tadalafil – used for the treatment of pulmonary arterial hypertension;
- St. John’s Wort (*Hypericum perforatum*) – an herbal product used to help with your mood.

Tell your doctor if you are taking any of the following medications as they may interact with boceprevir. The dosage of one or the other may have to be changed or the medication avoided:

- atorvastatin, bepridil, bosentan, budesonide, buprenorphine, colchicine, cyclosporine, desipramine, dexamethasone, digoxin, felodipine, fluticasone, fluvastatin, methadone, nicardipine, nifedipine, pravastatin, rifabutin, rosuvastatin, salmeterol, sirolimus, tacrolimus, trazodone, and warfarin.

How to use the peginterferon alfa-2b REDIPEN[®]



The following instructions explain how to inject yourself with the single use peginterferon alfa-2b REDIPEN[®]. **Please read the instructions carefully and completely before attempting to use the pen and follow them step by step.** Your doctor or his/her assistant will instruct you on how to self-inject with the peginterferon alfa-2b REDIPEN[®]. Do not attempt to inject yourself unless you are sure you understand the procedure and requirements for self-injection.

The peginterferon alfa-2b REDIPEN[®] is intended for use by one person only and must not be shared. Use the injection needle and alcohol swabs provided in the packaging only for the peginterferon alfa-2b REDIPEN[®]. Be sure the solution is at room temperature at the time of injection. Your doctor will have told you what dose you require for your therapy.

Note: The color of the dosing button is different for each strength of the peginterferon alfa-2b REDIPEN[®].

Step 1: Mixing

It is important that you keep the peginterferon alfa-2b REDIPEN[®] upright (as shown in Figure 1) during mixing, unless otherwise instructed.

- Take the peginterferon alfa-2b REDIPEN[®] out of the refrigerator. Allow the medicine to come to room temperature.
- Wash your hands with soap and water.

PROPER USE OF THIS MEDICATION

- Take VICTRELIS TRIPLE[™] exactly as your healthcare provider tells you. Your healthcare provider will tell you how much to take and when to take it.
- Tell your doctor, pharmacist or health professional if you notice any change in the appearance of boceprevir capsule, ribavirin capsule or peginterferon alfa-2a powder for solution.
- Always take VICTRELIS[™] and ribavirin with food.

The following instructions explain how to reconstitute and inject peginterferon alfa-2b powder for solution yourself. Please read the instructions carefully and follow them step by step. Your doctor or his/her assistant will instruct you how to self-inject peginterferon alfa-2b. Do not attempt to inject yourself unless you are sure you understand the procedure and requirement of self-injection.

Carefully follow the instructions provided.

- Place the peginterferon alfa-2b REDIPEN[®] upright in the holder of the tray provided in the pack (the dosing button will be on the **bottom**) (Figure 1).



Figure 1

- You may want to hold the REDIPEN[®] using the grip. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the REDIPEN[®] click. The two stoppers should come together.
- Wait for several seconds until the powder is completely dissolved.
- Gently turn the peginterferon alfa-2b REDIPEN[®] upside down twice. To avoid excessive foaming, do not shake.**
- Maintaining the peginterferon alfa-2b REDIPEN[®] upright, check the mixed peginterferon alfa-2b solution through the window. If there is still foam, wait until it settles. The solution should be clear and colorless. **Do not use the pen if the solution is discolored or irregular or contains any particles.**
- Keeping the peginterferon alfa-2b REDIPEN[®] upright in the holder provided in the packaging, disinfect the rubber membrane of the peginterferon alfa-2b REDIPEN[®] with one alcohol swab.
- Take the injection needle provided in the tray and remove its protective paper tab.
- Maintaining the peginterferon alfa-2b REDIPEN[®] upright in the holder,** gently push the injection needle onto the REDIPEN[®] and **screw it securely in place** (Figure 2).



Figure 2

Keep the peginterferon alfa-2b REDIPEN[®] pointed up.

- Do not take off the outer needle cap at this point.**
- You may see some liquid trickle out from under the cap, as the air has been expelled out of the pen.
- Wait about 5 seconds for this process to finish.
- Check through the window to be sure that the two stoppers are together. If they are not together, do not use this pen because you may not be able to dial your dose (Figure 3).**




Figure 3

Step 2: Setting the dose

- Remove the peginterferon alfa-2b REDIPEN[®] from the holder.
- Holding the peginterferon alfa-2b REDIPEN[®] firmly, pull the dosing button out as far as it will go, until you see a **dark ring** on the pen. The dosing button should be easy to pull out without excessive force being required (Figure 4).



Figure 4

Note: Do not push the dosing button back in at this time. You will push it in when you are ready to self-inject the peginterferon alfa-2b.

- Turn the dosing button until your prescribed dose is aligned with the dosing tab. The button should turn easily without excessive force being required (Figure 5).



Figure 5

Note: If you cannot easily pull out the dosing button or dial the dose, do not use excessive force and do not use this pen because it may not deliver the correct dose.

Step 3: Injecting the solution

- Select the injection site. Your doctor will have told you which sites to use (e.g., thigh or abdomen).

Note: Change your injection site each time.

- Clean the injection site skin with the second alcohol swab.

- Pull off the **outer needle cap** (Figure 6).




Figure 6

- There may be some liquid around the inner needle cap. This liquid is not part of your dose, this is extra. This is normal, as the air has been expelled out of the needle.
- Once the injection site is dry, pull off the **yellow inner needle cap** carefully exposing the injection needle.
- Hold the peginterferon alfa-2b REDIPEN[®] with your fingers wrapped around the pen body and your thumb on the dosing button (Figure 7).



Figure 7

- With your other hand, pinch a fold of loose skin.
- Insert the needle into the pinched skin at an angle of 45° to 90°.
- Press the dosing button down **slowly** and **firmly** until the button can no longer move.
- **Maintain pressure on the dosing button for an**

additional 5 seconds to ensure that you get the complete dose.

- Remove the needle from your skin.
- Press the injection site with a small bandage or sterile gauze if necessary for a few seconds.
- Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.
- Discard the peginterferon alfa-2b REDIPEN[®] with the needle safely in a closed rigid container.

Usual Adult Dose:

Your doctor has determined the correct dose of ribavirin and peginterferon alfa-2b based on your weight and the regimen (plan of treatment) that you are following for hepatitis C. Boceprevir 800 mg Three Times Daily will be used in combination with peginterferon alpha and ribavirin. Boceprevir will only be added to the treatment from the start of the fifth week. The total duration of your treatment will depend on the way in which your virus responds to treatment.

Your doctor may adjust your dose and length of time you take this treatment according to your response. Blood tests will be done regularly to help your doctor to know if it is working and if the dose needs to be changed.

If you have or develop severe kidney or liver problems, this treatment will be stopped. This treatment is not recommended for use in patients under the age of 18 years.

Based on the results of a clinical trial, the recommended dose of peginterferon alfa-2b is 1.5 mcg/kg/week in combination with ribavirin, which are dosed by patient weight. Peginterferon alfa-2b should be administered as subcutaneous injection once a week.

Ribavirin capsules are taken daily. Ribavirin capsules are to be administered orally, 800–1,400 mg, each day in two divided doses with food (morning and evening).

PATIENTS WHO HAVE NEVER BEEN TREATED

Recommended Dose

peginterferon alfa-2b: 1.5 mcg/kg/week

ribavirin: 800–1,400 mg daily based upon patient weight

boceprevir: 800 mg (four 200 mg capsules) three times daily (every 7 to 9 hours)

Dosing Recommendations^a

Patient Weight (kg)	peginterferon alfa-2b Powder for Solution		ribavirin Capsules	
	Weekly Dose (mcg/kg)	REDIPEN [®] Size (mcg/0.5 mL) ^b	Daily Dose (mg)	Number of Capsules (200 mg)
40 to 50	1.5	80	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
51 to 65	1.5	100	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
66 to 80	1.5	120	1,000	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
81 to 105	1.5	150	1,200	3 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
> 105	1.5	c	1,400	3 x 200 mg capsules A.M. 4 x 200 mg capsules P.M.

- a: The daily dose for ribavirin approximately falls within 13 ± 2 mg/kg/day.
- b: When reconstituted as instructed
- c: Should be calculated based on the body weight of an individual patient

PATIENTS WHO FAILED PRIOR TREATMENT

Recommended Dose

Peginterferon alfa-2b: 1.5 mcg/kg/week
 Ribavirin: 800–1,400 mg daily based upon patient weight
 boceprevir: 800 mg (four 200 mg capsules) three times daily (every 7 to 9 hours)

Dosing Recommendations^a

Patient Weight (kg)	peginterferon alfa-2b Powder for Solution		ribavirin Capsules	
	Weekly Dose (mcg/kg)	REDIPEN [®] Size (mcg/0.5 mL) ^b	Daily Dose (mg)	Number of Capsules (200 mg)
40 to 50	1.5	80	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
51 to 65	1.5	100	800	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
66 to 85	1.5	120	1,000	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
86 to 105	1.5	150	1,200	3 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
> 105	1.5	c	1,400	3 x 200 mg capsules A.M. 4 x 200 mg capsules P.M.

- a: The daily dose for ribavirin approximately falls within 13 ± 2 mg/kg/day.
- b: When reconstituted as instructed
- c: Should be calculated based on the body weight of an individual patient

It is important to follow your dosing schedule and your doctor’s instructions on how to take your medications. Take the medicine for as long as prescribed and do not exceed the recommended dosage.

Always take boceprevir with food (a meal or a light snack, such as a piece of fruit or crackers).

Take the ribavirin capsules by mouth with water and during your meal. Do not chew the capsules.

Peginterferon alfa-2b is given at a dose of 1.5 mcg/kg once a week. Peginterferon alfa-2b is for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection.

If you are injecting peginterferon alfa-2b yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive. Inject peginterferon alfa-2b on the same day each week. Injecting it at the same time of day each week will help you not to forget to take it. Take the dose as soon as you remember, then continue your treatment as usual. Do not take a double dose to make up for forgotten individual doses. Contact your doctor or pharmacist if needed.

Overdose:

The primary effects of overdose were an increased incidence and severity of adverse events reported at the therapeutic doses of VICTRELIS TRIPLE™.

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose of boceprevir and it is less than 2 hours before the next dose is due, the missed dose should be skipped. If you miss a dose and it is more than 2 hours before the next dose is due, take the missed dose with food and continue the normal dosing schedule. Do not double the next dose. If you have questions about what to do, call your healthcare provider.

If you miss a dose of ribavirin, take the missed dose as soon as possible during the same day. If an entire day has gone by, check with your doctor about what to do. Do not double the next dose.

If you miss a dose of peginterferon alfa-2b, take the missed dose as soon as possible during the same day or on the next day, and continue the dosing schedule provided to you by your doctor. If several days go by, check with your doctor about what to do. Do not double the next dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

What are the possible side effects of VICTRELIS TRIPLE™ therapy?

Like all medicines, VICTRELIS TRIPLE™ can have side effects. Although not all of these side effects may occur, they may need medical attention if they do occur.

VICTRELIS TRIPLE™ may cause serious side effects, including:

- Blood problems. Boceprevir can cause low red cell counts (anemia) or low neutrophil (neutropenia), which is a type of white blood cell, and low platelet counts (thrombocytopenia). In some people, these blood counts may fall to dangerously low levels.

Check with your doctor immediately if any of the following side effects occur during treatment:

- chest pain or persistent cough;
- symptoms of a severe allergic reaction (such as difficulty breathing, wheezing, or hives);
- symptoms associated with a cold or other respiratory infection, such as difficulty breathing or cough;
- shortness of breath;
- fever or chills beginning after a few weeks of treatment;
- changes in the way your heart beats;
- feeling depressed or hopeless, or thinking about death (suicidal thoughts or attempts);
- confusion, aggressiveness (sometimes directed against others), hallucination;
- trouble sleeping;
- severe stomach pain, black or tar-like stools, blood in stool or urine, feelings of numbness or tingling;
- severe bleeding from your nose;
- lower back or side pain, painful or difficult urination;
- problems with your eyesight or hearing;
- you notice that you are unusually tired and pale, and bruise easily.

Other events that may occur with this treatment are:

- irritation or pain at the site of injection;
- general discomfort, such as headache, fatigue or sleepiness, chills, fever, “flu-like” symptoms, weakness, pain around the ribs on the right side, feeling generally unwell, flushing, increased sweating;
- high or low blood pressure;
- dizziness, vertigo or faintness;
- sore tongue or mouth, dry mouth, thirst, loss of appetite, weight loss, nausea (feeling sick), vomiting, stomach or abdominal pain, indigestion, gas, diarrhea, loose stools, constipation;
- muscle ache, pain or stiffness, joint pain, arthritis;
- irritability, anxiety, agitation, nervousness, mood swings, difficulty concentrating, lack of interest in life;
- loss of hair or change in hair;
- skin disorders, including itching or rash, dry skin, redness, brown spots on skin, increased or decreased sensitivity to touch, sensitivity to light, eczema, psoriasis;
- disorders of the respiratory tract, including hoarseness, sore throat, cough, runny nose, stuffy nose, sinus infection, bronchitis, pneumonia;
- viral or fungal infection, herpes simplex (fever blister); or
- menstrual disorder.

The most common side effects of VICTRELIS TRIPLE™ include:

- fatigue, low red blood cell count (anemia), change in sense of taste, nausea, headache, diarrhea, vomiting, abdominal pain, fever, muscle and joint pain, weight loss, difficulty in sleeping, and dry skin;

- fatigue, dizziness, fainting, changes in blood pressure, and blurred vision can occur, so be cautious before driving or operating heavy machinery.

Some patients may have: change in sense of taste or smell, inflammation, infection, pain, or dryness of the eye, tear disorder, blurred vision, earache, middle ear infection, allergic reaction, puffiness of hands and feet, inflamed or bleeding gums, tooth abscess, rectal sores, decreased sex drive, impotence, irritation of the vagina, migraine headache, gout, change in thyroid function.

Very rarely, cases of stroke (cerebrovascular events) have been reported.

Very rarely, peginterferon alfa-2b, alone or with ribavirin, may cause aplastic anemia. Aplastic anemia is a condition caused by the failure of the bone marrow to make new red blood cells, white blood cells and platelets. Pure red cell aplasia has also been reported. Pure red cell aplasia is a condition in which severe and sudden anemia (characterized by symptoms such as severe tiredness/fatigue, and shortness of breath on mild exertion) develops due to failure of the bone marrow to produce red blood cells.

Additionally, the following events have been reported with peginterferon alfa-2b: facial palsy (weakness and slumping on one side to the face), severe allergic reactions such as angioedema (an allergic skin disease characterized by patches of circumscribed swelling involving the skin and its subcutaneous layers, the mucous membranes, and sometimes the internal organs), toxic epidermal necrolysis/Stevens Johnson Syndrome/erythema multiforme (a spectrum of rashes with varying degree of severity including death which may be associated with blisters in the mouth, nose, eyes and other mucosal membranes and sloughing of the affected area of the skin) and blindness.

Additionally, Vogt-Koyanagi-Harada syndrome (an autoimmune inflammatory disorder affecting the eyes, skin and the membranes of the ears, brain and spinal cord) has been reported with peginterferon alfa-2b use.

Tell your provider right away if you have any side effect that bothers you or that does not go away.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom/effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Very common	Blood problems: low red cell counts (anemia) which may lead to tiredness, headaches, shortness of breath when exercising, dizziness and looking pale.		√	
	Blood problems: low white blood cell counts (neutropenia) which may lead to an increased risk of getting infections.		√	
Common	Mental health: depression, thoughts of suicide, experience hallucinations, aggressiveness or confusion, or have trouble sleeping or concentrating		√	
	Heart: chest pain, high or low blood pressure, changes in the way your heart beats		√	
	Infection: High fever or chills, or pain while urinating		√	
	Thyroid: new or worsening problems with thyroid function		√	
Uncommon	Blood sugar: high blood sugar or diabetes		√	
	Colitis (inflammation of the bowel): abdominal pain, bloody diarrhea, fever		√	
	Eye: change in vision such as decrease or loss of vision		√	
	Ear: hearing problem		√	
	Lung: trouble breathing, infection, pneumonia, inflammation of lung tissue, new or worse high blood pressure in the lung (pulmonary hypertension)		√	
	New or worsening rheumatoid arthritis, systemic lupus erythematosus, psoriasis		√	
	Women who are planning or become pregnant			√

This is not a complete list of side effects. For any unexpected effects while taking VICTRELIS TRIPLE™, contact your doctor or pharmacist.

HOW TO STORE IT

Storage of VICTRELIS TRIPLE™ Packages:

Store the VICTRELIS TRIPLE™ package refrigerated between 2°C and 8°C.

Storage of boceprevir:

Boceprevir Capsules should be refrigerated at 2°C–8°C until dispensed by a pharmacist. When separated and for patient use, refrigerated capsules of VICTRELIS™ can remain stable until the expiration date printed on the label.

VICTRELIS™ can also be stored at room temperature (15°C–30°C) for 3 months. Store in the original container.

Storage of ribavirin:

When separated, ribavirin capsules should be stored in the refrigerator between 2°C and 8°C or at controlled room temperature between 15°C and 30°C.

Stability and storage for peginterferon alfa-2b REDIPEN®:

Store the peginterferon alfa-2b REDIPEN® at 2°C to 8°C. Once reconstituted peginterferon alfa-2b REDIPEN® should be used immediately but may be stored at 2°C–8°C for up to 24 hours. Do not freeze.

Keep VICTRELIS TRIPLE™ and all medicines out of the reach and sight of children.

Do not use past expiry date on the label.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701E
Ottawa, Ontario K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

or at Merck Canada Inc. by one of the following 2 ways:

- Call toll-free at 1-800-567-2594
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-496-9092, or
 - Mail to: Merck Canada Inc.
Pharmacovigilance
P.O. Box 1005
Pointe-Claire–Dorval, QC
H9R 4P8

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program or Merck does not provide medical advice.

MORE INFORMATION

For this document plus the full product monograph, prepared for health professionals, please contact the sponsor, Merck Canada Inc. at: 1-800-567-2594.

This leaflet was prepared by Merck Canada Inc.

™Trademarks of Schering Corporation, a subsidiary of **Merck & Co., Inc.** Used under license.

PEGETRON® is a registered trademark of Schering-Plough Ltd. Used under license.

REDIPEN® is a registered trademark of Merck Canada Inc.

* All other trademarks are the property of their respective owner(s).

© 2011, Merck Canada Inc., a subsidiary of **Merck & Co., Inc.** All rights reserved.

Last revised: November 4, 2011

